arXiv:1404.6432v1 [math.NT] 25 Apr 2014

Modular case of Levinson's theorem

BERNARD Damien

Abstract

We evaluate the integral mollified second moment of L-functions of primitive cusp forms and we obtain, for such L-function, an explicit positive proportion of zeros which lie on the critical line.

Contents

1 Introduction and overview of the results **1**

2 Review on L-functions of primitive cusp forms **4**

3 Mollified second moment of L-functions of modular forms **5**
3.1 Evaluation of the off-diagonal term 7
3.1.1 Initial lemmas 8
3.1.2 Shifted convolution sums 10
3.1.3 Proof of proposition 1 16
3.2 Evaluation of the diagonal term 16
3.2.1 Initial lemmas 17
3.2.2 Estimation of $I_f^{D_1}(\alpha,\beta)$ 20
3.2.3 Proof of proposition 4 26

4 Effective proportion of zeros on the critical line **27**

5 Non-mollified second integral moment **28**
5.1 Diagonal contribution 29
5.2 Conjecture of Conrey, Farmer, Keating, Rubinstein and Snaith 31

1 Introduction and overview of the results

Nowadays, we know that more than 41% of the non-trivial zeros of the Riemann zeta-function lie on the critical line ([BCY11] and [Fen12]). This theorem is the best one of a sequence of results about the percentage of zeros ρ satisfying $\Re\rho = 1/2$.

Historically, Selberg ([Sel42]) was the first one to show that this proportion is not zero without quantifying it. According to Titchmarsh ([Tit86], part 10.9), it was calculated later on in Min's dissertation, that the proportion obtained by Selberg's method is very small. One may

refer to the introduction of [Ste07] (page 8) for numerical values. In 1974, Levinson ([Lev74]) succeeded in proving that at least one third of the non-trivial zeros lie on the critical line by perturbing the Riemann zeta function by a linear combination of its derivatives. A significant improvement, due to Conrey ([Con89]), increased this proportion up to more than two fifths. In order to do this, he improves the general result of [BCHB85] on the asymptotic behaviour of the mollified second moment of the Riemann zeta function, when the coefficients of the mollifier are essentially given by the Moebius function, which allows him to work with a longer mollifier than Levinson's one. From this last result of Conrey and using a two parts mollifier, Bui, Conrey and Young ([BCY11]) proved that 41% of non-trivial zeros ρ satisfies $\Re\rho = 1/2$.

Since the Riemann zeta-function is a L-function of degree one, it is rather natural to generalise these results to L-functions of higher degrees. For instance, Hafner has extended Selberg's result to L-functions of degree two. Precisely, if f is a holomorphic cusp form of even weight and full level or an even Maaß form of full level, let $N_f(T)$ (resp. $N_{f,0}(T)$) be the number of the non-trivial zeros ρ (resp. on the critical line) of $L(f,s)$ with $0 < \Im(\rho) \leq T$. Then, Hafner proved in [Haf83] and [Haf87] that there exists a positive number A such that $N_{f,0}(T) > AN_f(T)$ for large T. Rezvyakova ([Rez10]) adapted ([Haf83]) to L-functions attached to automorphic cusp forms for congruence subgroups. Nevertheless, they do not give any explicit value for A but by analogy with the Riemann zeta-function case, this constant should probably be close to zero.

In [Far94], Farmer applied Levinson's method to L-functions of an holomorphic cusp form f of even weight and full level and succeeded in determining the asymptotic behaviour of the mollified integral second moment of $L(f,s)$ when the mollifier is a Dirichlet polynomial of length less than $T^{1/6-\varepsilon}$. From this result, Farmer obtained explicit lower bounds for the proportion of simple zeros of the j^{th} derivative ($j \geq 1$) of the completed L-function of $L(f,s)$, which are on the critical line. Unfortunately, the length of the mollifier is too small to exhibit an explicit positive proportion of simple zeros on the critical line for $L(f,s)$ itself. Nevertheless, even thought Farmer did not remark it, his result proves that at least $1,65\%$ of zeros of $L(f,s)$ satisfy $\Re s = 1/2$ (confer part 4).

In this paper, we exhibit a positive proportion of zeros, which lie on the critical line for L-functions of holomorphic primitive cusp forms. To get this result, we study the asymptotic behaviour of the smooth mollified second moment of $L(f,s)$ following the method developed in [You10]. We choose a mollifier ψ, which is a Dirichlet polynomial of length T^ν defined in relation (7) page 5. Moreover, we introduce a smooth function w compactly supported in $[T/4, 2T]$ with some conditions on its derivatives (confer relations (8a), (8b) and (8c)). We prove the following theorem.

Theorem 1 *Let f be a holomorphic primitive cusp form of even weight, square-free level and trivial character. If $0 < \nu < \frac{1-2\vartheta}{4+2\vartheta}$ and if α, β are complex numbers satisfying $\alpha, \beta \ll L^{-1}$ with $|\alpha+\beta| \gg L^{-1}$, then*

$$\int_{-\infty}^{+\infty} w(t) L\left(f, \frac{1}{2}+\alpha+it\right) L\left(f, \frac{1}{2}+\beta-it\right) |\psi(\sigma_0+it)|^2\, dt = \widehat{w}(0)c(\alpha,\beta) + O(T(\ln L)^4/L)$$

where

$$c(\alpha,\beta) = 1 + \frac{1}{\nu}\frac{1-T^{-2(\alpha+\beta)}}{(\alpha+\beta)\ln T}\ \frac{d^2}{dxdy}\left[M^{-\beta x-\alpha y}\int_0^1 P(x+u)P(y+u)du\right]\Bigg|_{x=y=0} \tag{1}$$

and where $\theta = 7/64$ refers to the exponent in the approximation towards Ramanujuan-Petersson-Selberg conjecture (confer relation (16)*).*

Corollary 1 *Let f be a holomorphic primitive cusp form of even weight, square-free level and trivial character. At least $2,97\%$ of non-trivial zeros of $L(f,s)$ lie on the critical line $\Re(s) = 1/2$. Assuming the Selberg conjecture, we improve this percentage to $6,93\%$. In other words,*

$$\liminf_{T\to+\infty} \frac{N_{f,0}(T)}{N_f(T)} \geq \begin{cases} 0,0297 & \textit{unconditionally,} \\ 0,0693 & \textit{under Selberg conjecture.} \end{cases}$$

Remark 1 As Heath-Brown ([HB79]) and Selberg pointed out, the study of the second mollified moment allows to obtain a lower bound for the proportion of the simple non-trivial zeros lying on the critical line. Unfortunately, in our case, the length of the mollifier is too small to get a positive proportion of simple zeros satisfying the Riemann hypothesis for now. We plan to get back to this issue in the close future.

The method we used can also be applied to determine the asymptotic behaviour of the smooth second moment of $L(f,s)$ close to the critical line.

Theorem 2 *Let f be a holomorphic primitive cusp form of even weight, square-free level N and trivial character. If α, β are complex numbers satisfying $\alpha, \beta \ll L^{-1}$, we have*

$$\int_{-\infty}^{+\infty} w(t) L\left(f, \frac{1}{2}+\alpha+it\right) L\left(f, \frac{1}{2}+\beta-it\right) dt = \mathfrak{a}_f \int_{-\infty}^{+\infty} w(t) \ln t dt + \left[\mathfrak{b}_f + \mathfrak{a}_f \ln\left(\frac{\sqrt{N}}{2\pi}\right)\right] \widehat{w}(0) + O\left(|\alpha+\beta| T (\ln T)^2 + T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

with

$$\mathfrak{a}_f = \frac{12N}{\pi^2 \nu(N)} L\left(Sym^2 f, 1\right) \text{ and } \mathfrak{b}_f = \frac{12N}{\pi^2 \nu(N)} L\left(Sym^2 f, 1\right) \left(\frac{L'\left(Sym^2 f, 1\right)}{L\left(Sym^2 f, 1\right)} + \gamma + \sum_{p|N} \frac{\ln p}{p+1} - \frac{2\zeta'(2)}{\zeta(2)} \right)$$

and where $\nu(N) = N \prod_{p|N} \left(1 + \frac{1}{p}\right)$.

Remark 2 Our result is non-trivial only in the case $|\alpha+\beta| = o(1/\ln T)$, and furthermore, we need $|\alpha+\beta| = o(1/\ln^2 T)$ to ensure that the term of order T is significant.

Corollary 2 *Let f be a holomorphic primitive cusp form of even weight, square-free level N and trivial character. Then*

$$\int_{-\infty}^{+\infty} w(t) \left| L\left(f, \frac{1}{2}+it\right)\right|^2 dt = \int_{-\infty}^{+\infty} w(t) \left[\mathfrak{a}_f \ln\left(\frac{t\sqrt{N}}{2\pi}\right) + \mathfrak{b}_f\right] dt + O\left(T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

Remark 3 This corollary is in agreement with the conjecture stated in [CFK$^+$05] about integral moments of L-functions (see section 5.2).

In [Zha05], Zhang succeeded in determining the main term of this integral second moment of $L(f,s)$ (without smooth function w) on the critical line. Thanks to corollary 2, we improve his result with the following more precise asymptotic expansion. When f is a holomorphic cusp form of even weight for the full modular group, we can also refer to [Goo82] where a similar asymptotic expansion is given.

Corollary 3 *Let f be a holomorphic primitive cusp form of even weight, square-free level N and trivial character. We have*

$$\int_0^T \left| L\left(f, \frac{1}{2}+it\right)\right|^2 dt = \mathfrak{a}_f T\ln T + \left[\mathfrak{b}_f + \mathfrak{a}_f \ln\left(\frac{\sqrt{N}}{2\pi e}\right)\right] T + O\left(T/\ln T\right).$$

Notations

- If f and g are some functions of the real variable, $f(x) \ll_A g(x)$ or $f(x) = O_A(g(x))$ mean that $|f(x)|$ is smaller than a constant, which only depends on A, times $|g(x)|$ for large x.
- Similarly, the notation $f \asymp g$ means $f(x) \ll g(x)$ and $g(x) \ll f(x)$.
- The function ln refers to the natural logarithm function.
- From the Riemann zeta function ζ, we define for any positive square-free integer N

$$\zeta^{(N)}(s) = \prod_{p|N}\left(1-\frac{1}{p^s}\right)\zeta(s).$$

Acknowledgements I would like to thank Guillaume Ricotta and Emmanuel Royer for all their comments and their remarks during this work. I want to express my gratitude to them for their kindness and for all their encouragements. I wish to thank the organisers of the Number Theory Seminar of Institut de Mathématiques de Bordeaux for their warm welcome in may 2012.

2 Review on L-functions of primitive cusp forms

For this section, we may refer to [IK04] (chapter 14). Throughout this paper, f denotes a holomorphic primitive cusp form of even weight k and square-free level N. The Fourier expansion at the cusp ∞ of f is given by

$$f(z) = \sum_{n\geq 1} \lambda_f(n) n^{\frac{k-1}{2}} e^{2i\pi nz}$$

for every complex number z in the upper-half plane with the arithmetic normalisation $\lambda_f(1)=1$. The Fourier coefficients $\lambda_f(n)$ satisfy the multiplicative relations

$$\lambda_f(n)\lambda_f(m) = \sum_{\substack{d|(m,n)\\(d,N)=1}} \lambda_f\left(\frac{mn}{d^2}\right) \tag{2}$$

$$\lambda_f(mn) = \sum_{\substack{d|(m,n)\\(d,N)=1}} \mu(d)\lambda_f\left(\frac{m}{d}\right)\lambda_f\left(\frac{n}{d}\right) \tag{3}$$

for all positive integers m and n. Since $\lambda_f(1) \neq 0$, we may define the convolution inverse $(\mu_f(n))$ of the sequence $(\lambda_f(n))$. This is an arithmetic multiplicative function, which satisfies for a prime number p

$$\mu_f(1) = 1, \quad \mu_f(p) = -\lambda_f(p), \quad \mu_f(p^2) = \lambda_f(p)^2 - \lambda_f(p^2) = \begin{cases} 1 & \text{if } p \nmid N \\ 0 & \text{otherwise} \end{cases} \tag{4}$$

$$\text{and if } j \geq 3 \quad \mu_f(p^j) = 0. \tag{5}$$

We consider

$$L(f,s) = \sum_{n\geq 1} \frac{\lambda_f(n)}{n^s} = \prod_{p\in\mathcal{P}} \left(1 - \frac{\lambda_f(p)}{p^s} + \chi_0(p)\frac{1}{p^{2s}}\right)^{-1} = \prod_{p\in\mathcal{P}} \left(1 - \frac{\alpha_f(p)}{p^s}\right)^{-1} \left(1 - \frac{\beta_f(p)}{p^s}\right)^{-1},$$

which is an absolutely convergent and non-vanishing Dirichlet series, an Euler product on $\Re(s) > 1$, where χ_0 denotes the trivial character of modulus N and $\alpha_f(p)$, $\beta_f(p)$ are the complex roots of the quadratic equation $X^2 - \lambda_f(p)X + \chi_0(p) = 0$. Moreover, the function

$$\Lambda(f,s) = \left(\frac{\sqrt{N}}{2\pi}\right)^s \Gamma\left(s + \frac{k-1}{2}\right) L(f,s) = L_\infty(f,s)L(f,s)$$

is the completed L-function of $L(f,s)$. It can be extended to an holomorphic function on $\mathbb{C}$ and satisfies the functional equation

$$\Lambda(f,s) = \varepsilon(f)\Lambda(f,1-s)$$

where $\varepsilon(f) = \pm 1$. Remark that, by the duplication formula for the gamma function, the local factor at infinity can be written

$$L_\infty(f,s) = \left(\frac{2^k}{8\pi}\right)^{1/2} \left(\frac{\sqrt{N}}{\pi}\right)^s \Gamma\left(\frac{s}{2} + \frac{k-1}{4}\right) \Gamma\left(\frac{s}{2} + \frac{k+1}{4}\right). \tag{6}$$

3 Mollified second moment of L-functions of modular forms

This section contains the proof of theorem 1. We define the sequence $(\mu_f(n))_{n\geq 1}$ as the convolution inverse of the sequence $(\lambda_f(n))_{n\geq 1}$ and we define a mollifier ψ of the shape

$$\psi(s) = \sum_{n\leq M} \frac{\mu_f(n)}{n^{s+\frac{1}{2}-\sigma_0}} P\left(\frac{\ln M/n}{\ln M}\right) \tag{7}$$

with $M = T^\nu$, $\sigma_0 = \frac{1}{2} - \frac{R}{\ln T}$ where R is a positive real number and P is a real polynomial satisfying $P(0) = 0$, $P(1) = 1$. In addition, we choose a function $w : \mathbb{R} \mapsto \mathbb{R}$, which satisfies

$$w \text{ is smooth,} \tag{8a}$$

$$w \text{ is compactly supported with supp } w \subset [T/4; 2T], \tag{8b}$$

$$\text{for each } j \geq 0, \text{ we have } w^{(j)}(t) \ll_j \Delta^{-j} \text{ where } \Delta = T/L \text{ and } L = \ln T. \tag{8c}$$

For more convenience, we set

$$I_f(\alpha,\beta)=\int_{-\infty}^{+\infty} w(t)L\left(f,\frac{1}{2}+\alpha+it\right)L\left(f,\frac{1}{2}+\beta-it\right)|\psi(\sigma_0+it)|^2\,dt.$$

To study the asymptotic behaviour of $I_f(\alpha,\beta)$, we need an explicit expression of $L(f,s)$ with $0\leq\Re(s)\leq 1$. In lemma 1, we get an exact formula, also called "approximate functional equation", which gives an expression for $L(f,s+it)L(f,s-it)$ with s in the critical strip where we can not use the Dirichlet series. Thanks to this new relation, we may split $I_f(\alpha,\beta)$ in a diagonal term (without oscillation) and an off-diagonal term (with oscillation). The off-diagonal contribution is bounded in part 3.1 whereas the diagonal term is estimated in part 3.2.

Lemma 1 *Let G be any entire function, which decays rapidly in vertical strips, even and normalised by $G(0)=1$. Then for each complex numbers α, β such that $0\leq|\Re(\alpha)|,|\Re(\beta)|\leq 1/2$, we have*

$$L\left(f,\frac{1}{2}+\alpha+it\right)L\left(f,\frac{1}{2}+\beta-it\right)=\sum_{m,n\geq 1}\sum\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha}n^{\frac{1}{2}+\beta}}\left(\frac{m}{n}\right)^{-it}V_{\alpha,\beta}(mn,t)$$
$$+X_{\alpha,\beta,t}\sum_{m,n\geq 1}\sum\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}-\beta}n^{\frac{1}{2}-\alpha}}\left(\frac{m}{n}\right)^{-it}V_{-\beta,-\alpha}\left(mn,t\right)$$

where

$$g_{\alpha,\beta}(s,t)=\frac{L_\infty\left(f,\frac{1}{2}+\alpha+s+it\right)L_\infty\left(f,\frac{1}{2}+\beta+s-it\right)}{L_\infty\left(f,\frac{1}{2}+\alpha+it\right)L_\infty\left(f,\frac{1}{2}+\beta-it\right)},\ V_{\alpha,\beta}(x,t)=\frac{1}{2i\pi}\int_{(1)}\frac{G(s)}{s}g_{\alpha,\beta}(s,t)x^{-s}ds$$

and

$$X_{\alpha,\beta,t}=\frac{L_\infty\left(f,\frac{1}{2}-\alpha-it\right)L_\infty\left(f,\frac{1}{2}-\beta+it\right)}{L_\infty\left(f,\frac{1}{2}+\alpha+it\right)L_\infty\left(f,\frac{1}{2}+\beta-it\right)}.$$

We do not write the proof of this lemma, which is essentially the same as theorem 5.3 of [IK04]. Nevertheless, it will be usefull to have good approximations of $X_{\alpha,\beta,t}$, $g_{\alpha,\beta}(s,t)$ and $V_{\alpha,\beta}(x,t)$.

Lemma 2 *For large t and for $s\ll t^\varepsilon$ in any vertical strip, we have*

$$X_{\alpha,\beta,t} = \left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}\left(1+\frac{i(\alpha^2-\beta^2)}{t}+O\left(\frac{1}{t^2}\right)\right) \tag{9}$$

$$\text{and } g_{\alpha,\beta}(s,t) = \left(\frac{t\sqrt{N}}{2\pi}\right)^{2s}\left(1+O\left(\frac{|s^2|}{t}\right)\right). \tag{10}$$

In addition, for each integer $j\geq 0$ and for all real number $A>0$, we have

$$t^j\frac{\partial^j}{\partial t^j}V_{\alpha,\beta}(x,t)\ll_{A,j}\left(1+\frac{|x|}{t^2}\right)^{-A}. \tag{11}$$

Proof: We may write

$$\begin{aligned} X_{\alpha,\beta,t} &= \left(\frac{\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)} \frac{\Gamma\left(\frac{k}{2}-\alpha-it\right)\Gamma\left(\frac{k}{2}-\beta+it\right)}{\Gamma\left(\frac{k}{2}+\alpha+it\right)\Gamma\left(\frac{k}{2}+\beta-it\right)} \\ \text{and } g_{\alpha,\beta}(s,t) &= \left(\frac{\sqrt{N}}{2\pi}\right)^{2s} \frac{\Gamma\left(\frac{k}{2}+\alpha+s+it\right)\Gamma\left(\frac{k}{2}+\beta+s-it\right)}{\Gamma\left(\frac{k}{2}+\alpha+it\right)\Gamma\left(\frac{k}{2}+\beta-it\right)}. \end{aligned}$$

Then, the first part of this lemma comes from the following Stirling formula with $s=\sigma+i\tau$ in any vertical strip

$$\Gamma(s)=\sqrt{2\pi}|\tau|^{\sigma-\frac{1}{2}}e^{-\frac{\pi}{2}|\tau|}e^{i\left(\tau\ln|\tau|-\tau+\frac{\pi}{2}(\sigma-1/2)sgn(\tau)\right)}\left(1-i\frac{\left(\sigma-\frac{1}{2}\right)^2-\frac{1}{12}}{2\tau}+O\left(\frac{1}{\tau^2}\right)\right).$$

We refer to ([Ten95], corollaire 0.13). To prove (11), we move the integration line far to the right on $\Re s=A$ and by (10), we obtain the desired bound if $t^2\ll x$. In the case $x\ll t^2$, the result comes easily by trivial bounds.

□

Thanks to the previous fonctional equation, we may split $I_f(\alpha,\beta)$ as a sum of diagonal terms and off-diagonal terms. Precisely, opening the mollifier ψ, we may write

$$\begin{aligned} I_f(\alpha,\beta) &= \sum_{a,b\leq M}\frac{\mu_f(a)\mu_f(b)}{\sqrt{ab}}P\left(\frac{\ln M/a}{\ln M}\right)P\left(\frac{\ln M/b}{\ln M}\right)\Big[I_{a,b}^{D_1}(\alpha,\beta)+I_{a,b}^{D_2}(\alpha,\beta) \\ &\qquad +I_{a,b}^{ND_1}(\alpha,\beta)+I_{a,b}^{ND_2}(\alpha,\beta)\Big] \end{aligned} \tag{12}$$

with

$$\begin{aligned} I_{a,b}^{D_1}(\alpha,\beta) &= \sum_{am=bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha}n^{\frac{1}{2}+\beta}}\int_{-\infty}^{+\infty}w(t)V_{\alpha,\beta}(mn,t)dt, \\ I_{a,b}^{D_2}(\alpha,\beta) &= \sum_{am=bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}-\beta}n^{\frac{1}{2}-\alpha}}\int_{-\infty}^{+\infty}w(t)X_{\alpha,\beta,t}V_{-\beta,-\alpha}(mn,t)dt, \\ I_{a,b}^{ND_1}(\alpha,\beta) &= \sum_{am\neq bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha}n^{\frac{1}{2}+\beta}}\int_{-\infty}^{+\infty}w(t)\left(\frac{am}{bn}\right)^{-it}V_{\alpha,\beta}(mn,t)dt, \\ I_{a,b}^{ND_2}(\alpha,\beta) &= \sum_{am\neq bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}-\beta}n^{\frac{1}{2}-\alpha}}\int_{-\infty}^{+\infty}w(t)X_{\alpha,\beta,t}\left(\frac{am}{bn}\right)^{-it}V_{-\beta,-\alpha}(mn,t)dt. \end{aligned}$$

3.1 Evaluation of the off-diagonal term

In this part, we evaluate the size of the off-diagonal term. Precisely, we prove the following proposition.

Proposition 1 *If $0 < \nu < \frac{1-2\theta}{4+2\theta}$ and if α, β are complex numbers satisfying $\alpha, \beta \ll L^{-1}$ then there exists $\varepsilon > 0$ such that*

$$\sum_{a,b\leq M} \frac{\mu_f(a)\mu_f(b)}{\sqrt{ab}} P\left(\frac{\ln M/a}{\ln M}\right) P\left(\frac{\ln M/b}{\ln M}\right) \left[I_{a,b}^{ND_1}(\alpha,\beta) + I_{a,b}^{ND_2}(\alpha,\beta)\right] \ll T^{1-\varepsilon}.$$

The main tool of the proof of this proposition is a theorem about shifted convolution sums on average.

3.1.1 Initial lemmas

In order to prove the previous proposition, we begin to get rid of some harmless terms occurring in the definition of $I_{a,b}^{ND_1}(\alpha,\beta)$.

Lemma 3 *Let $\varepsilon > 0$, $0 < \gamma < 1$, $\alpha, \beta \ll L^{-1}$ be complex numbers and $a, b \leq T^{\nu}$ be positive integers. Then, for all real number $A > 0$, we have*

$$I_{a,b}^{ND_1}(\alpha,\beta) = \sum_{\substack{am\neq bn \\ mn\ll T^{2+\varepsilon} \\ \left|\frac{am}{bn}-1\right|\ll T^{-\gamma}}} \frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha} n^{\frac{1}{2}+\beta}} \int_{-\infty}^{+\infty} w(t) \left(\frac{am}{bn}\right)^{-it} V_{\alpha,\beta}(mn,t)dt + O\left(T^{-A}\right). \tag{13}$$

Proof: Firstly, by (11) with $j = 0$, we get for all real number $A > 0$

$$\int_{-\infty}^{+\infty} w(t) \left(\frac{am}{bn}\right)^{-it} V_{\alpha,\beta}(mn,t)dt \ll_A T \left(\frac{T^2}{mn}\right)^A.$$

As a consequence, for $A > 1/2$, since $|\lambda_f(n)| \leq \tau(n) \ll n^{\epsilon}$ and $\alpha, \beta \ll L^{-1}$, we may write

$$\begin{aligned}
\sum_{\substack{am\neq bn \\ mn>T^{2+\varepsilon}}} \frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha} n^{\frac{1}{2}+\beta}} \int_{-\infty}^{+\infty} w(t) \left(\frac{am}{bn}\right)^{-it} V_{\alpha,\beta}(mn,t)dt &\ll T^{1+2A} \sum_{mn>T^{2+\varepsilon}} \frac{\tau(m)\tau(n)}{m^{\frac{1}{2}+A+\Re\alpha} n^{\frac{1}{2}+A+\Re\beta}} \\
&\ll T^{1+2A} \sum_{h>T^{2+\varepsilon}} \frac{1}{h^{\frac{1}{2}+A-\epsilon}} \\
&\ll T^{2-A\varepsilon} \\
&\ll T^{-A}.
\end{aligned}$$

Then, using (11) for each integer j and since $w^{(j)}(t) \ll \Delta^{-j}$, for all real number $A > 0$, we have, uniformly with respect to x, the following bound

$$\frac{\partial^j}{\partial t^j}\left[w(t)V_{\alpha,\beta}(x,t)\right] \ll \Delta^{-j} \left(\frac{T^2}{|x|}\right)^A.$$

Hence, if $am \neq bn$, thanks to j integrations by parts, we get

$$\begin{aligned}
\int_{-\infty}^{+\infty} w(t) \left(\frac{am}{bn}\right)^{-it} V_{\alpha,\beta}(mn,t)dt &= \frac{1}{\left(i\ln\frac{am}{bn}\right)^j} \int_{-\infty}^{+\infty} \left(\frac{am}{bn}\right)^{-it} \frac{\partial^j}{\partial t^j}\left[w(t)V_{\alpha,\beta}(x,t)\right] dt \\
&\ll \frac{T}{\Delta^j \left|\ln\frac{am}{bn}\right|^j} \left(\frac{T^2}{mn}\right)^A.
\end{aligned}$$

Therefore, with $A = 1/2 + \max\{\Re\alpha, \Re\beta\} + \delta$ and $\delta > 0$, using the lower bound $x/2 \leq \ln(1+x)$ for $0 < x < 1$, we have

$$\sum_{\substack{am \neq bn \\ \left|\frac{am}{bn}-1\right| > T^{-\gamma}}} \frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha} n^{\frac{1}{2}+\beta}} \int_{-\infty}^{+\infty} w(t) \left(\frac{am}{bn}\right)^{-it} V_{\alpha,\beta}(mn,t)dt \ll \frac{T^{1+2A+j\gamma}}{\Delta^j} \sum_{m,n\geq 1} \frac{\tau(m)\tau(n)}{m^{\frac{1}{2}+\Re\alpha+A} n^{\frac{1}{2}+\Re\beta+A}}.$$

Since $\gamma < 1$, the result comes easily from a choice of large j.

□

We introduce a dyadic partition of unity to the sums over m and n. We fix an arbitrary smooth function $\rho :]0, +\infty[\to \mathbb{R}$, which is compactly supported in $[1, 2]$ and which satisfies

$$\sum_{\ell=-\infty}^{+\infty} \rho\left(2^{-\ell/2}x\right) = 1.$$

We may refer to [Har03] (section 5) to build such a function. For each integer ℓ, we define

$$\rho_\ell(x) = \rho\left(\frac{x}{A_\ell}\right) \text{ with } A_\ell = 2^{\ell/2}T^\gamma.$$

In order to study the asymptotic behaviour of $I_{a,b}^{ND_1}(\alpha,\beta)$, we consider the function $F_{h;\ell_1,\ell_2}$ which is defined by

$$F_{h;\ell_1,\ell_2}(x,y) = \frac{a^{\frac{1}{2}+\alpha} b^{\frac{1}{2}+\beta}}{x^{\frac{1}{2}+\alpha} y^{\frac{1}{2}+\beta}} \int_{-\infty}^{+\infty} w(t) \left(1+\frac{h}{y}\right)^{-it} V_{\alpha,\beta}\left(\frac{xy}{ab}, t\right) dt \times \rho_{\ell_1}(x)\rho_{\ell_2}(y). \tag{14}$$

Lemma 4 *Let $\varepsilon > 0$, $0 < \gamma < 1$, $\alpha, \beta \ll L^{-1}$ be complex numbers and $a, b \leq T^\nu$ be positive integers. Then, for all real number $A > 0$, we have*

$$I_{a,b}^{ND_1}(\alpha,\beta) = \sum_{\substack{A_{\ell_1}A_{\ell_2} \ll abT^{2+\varepsilon} \\ A_{\ell_1} \asymp A_{\ell_2} \\ A_{\ell_1}, A_{\ell_2} \gg T^\gamma}} \sum_{0<|h|\ll T^{-\gamma}\sqrt{A_{\ell_1}A_{\ell_2}}} \sum_{am-bn=h} \lambda_f(m)\lambda_f(n) F_{h;\ell_1,\ell_2}(am,bn) + O\left(T^{-A}\right). \tag{15}$$

Proof: For more convenience, we define

$$H(x,y) = \frac{a^{\frac{1}{2}+\alpha} b^{\frac{1}{2}+\beta}}{x^{\frac{1}{2}+\alpha} y^{\frac{1}{2}+\beta}} \int_{-\infty}^{+\infty} w(t) \left(\frac{x}{y}\right)^{-it} V_{\alpha,\beta}\left(\frac{xy}{ab}, t\right) dt.$$

From the previous lemma and using the partition of unity, we may write

$$I_{a,b}^{ND_1}(\alpha,\beta) = \sum_{\ell_1,\ell_2} \sum_{h\neq 0} \sum_{\substack{am-bn=h \\ mn \ll T^{2+\varepsilon} \\ \left|\frac{am}{bn}-1\right| \ll T^{-\gamma}}} \lambda_f(m)\lambda_f(n) H(am,bn)\rho_{\ell_1}(am)\rho_{\ell_2}(bn) + O\left(T^{-A}\right).$$

First, if $|h| \geq \sqrt{A_{\ell_1}A_{\ell_2}}T^{-\gamma}$ then

$$\max\left\{\left|\frac{am}{bn}-1\right|, \left|\frac{bn}{am}-1\right|\right\}^2 \geq \left|\frac{am}{bn}-1\right| \left|\frac{bn}{am}-1\right| = \frac{h^2}{ambn} \asymp \frac{h^2}{A_{\ell_1}A_{\ell_2}} \geq T^{-2\gamma}.$$

Secondly, if $|\ell_1 - \ell_2| \geq 3$, for instance if $\ell_1 - \ell_2 \geq 3$ then

$$\frac{am}{bn} - 1 \geq \frac{2^{\frac{\ell_1-\ell_2}{2}}}{2} - 1 \geq \sqrt{2} - 1 \gg 1.$$

Thus we may assume $A_{\ell_1} \asymp A_{\ell_2}$. Thirdly, if $A_{\ell_2} \leq T^{\gamma}$ then

$$\left|\frac{am}{bn} - 1\right| = \frac{h}{bn} \geq \frac{h}{2A_{\ell_2}} \gg T^{-\gamma}.$$

Therefore, we may assume $A_{\ell_2} \geq T^{\gamma}$ and, in the same way, $A_{\ell_1} \geq T^{\gamma}$. Finally, since $am - bn = h$, we get $H(am, bn)\rho_{\ell_1}(am)\rho_{\ell_2}(bn) = F_{h;\ell_1,\ell_2}(am, bn)$.

□

3.1.2 Shifted convolution sums

The core of the proof of our theorem is the following bound, which is a generalisation of theorem 2 of [Blo05], about shifted convolution sums on average. We define θ as the exponent in the Ramanujuan-Petersson conjecture, which claims

$$|\lambda(n)| \leq \tau(n)n^{\theta} \tag{16}$$

for eigenvalues $\lambda(n)$ of the Hecke operator T_n acting on the space of weight 0 Maaß cusp forms of level N.

Theorem 3 *Let ℓ_1, ℓ_2, H and h_1 be positive integers. Let M_1, M_2, P_1, P_2 be real numbers greater than 1. Let $\{g_h\}$ be a family of smooth functions, supported on $[M_1, 2M_1] \times [M_2, 2M_2]$ such that $||g_h^{(ij)}||_{\infty} \ll_{i,j} (P_1/M_1)^i(P_2/M_2)^j$ for all $i, j \geq 0$. Let $(a(h))$ be a sequence of complex numbers such that*

$$a(h) \neq 0 \Rightarrow h \leq H,\ h_1|h \text{ et } \left(h_1, \frac{h}{h_1}\right) = 1.$$

If $\ell_1 M_1 \asymp \ell_2 M_2 \asymp A$ and if there exists $\epsilon > 0$ such that

$$H \ll \frac{A}{\max\{P_1, P_2\}^2} \frac{1}{(\ell_1\ell_2 M_1 M_2 P_1 P_2)^{\epsilon}}, \tag{17}$$

then, for all real number $\varepsilon > 0$, we have

$$\sum_{h=1}^{H} a(h) \sum_{\substack{m_1,m_2 \geq 1 \\ \ell_1 m_1 - \ell_2 m_2 = h}} \lambda_f(m_1)\overline{\lambda_f(m_2)}g_h(m_1, m_2)$$

$$\ll A^{\frac{1}{2}} h_1^{\theta} \|a\|_2 (P_1 + P_2)^{\frac{3}{2}} \left[\sqrt{P_1 + P_2} + \left(\frac{A}{\max\{P_1, P_2\}}\right)^{\theta} \left(1 + \sqrt{\frac{(h_1, \ell_1\ell_2)H}{h_1\ell_1\ell_2}}\right)\right] (\ell_1\ell_2 M_1 M_2 P_1 P_2 H)^{\varepsilon}.$$

Proof: This proof is a direct generalisation of the proof of theorem 2 in [Blo05], that is why we only give an outline of the proof of Blomer. In order to simplify, we set

$$\Sigma(\ell_1,\ell_2,H,a)=\sum_{h=1}^{H}a(h)\sum_{\substack{m_1,m_2\geq 1\\ \ell_1 m_1-\ell_2 m_2=h}}\lambda_f(m_1)\overline{\lambda_f(m_2)}g_h(m_1,m_2).$$

We set also $\Delta=\min\left\{\frac{P_1}{\ell_1 M_1},\frac{P_2}{\ell_2 M_2}\right\}$, let $Q\geq 1/\Delta$ be a very large parameter and set $\delta=1/Q$ such that $\delta\leq\Delta$. Let ϕ be a smooth function compactly supported in $[-\Delta^{-1},\Delta^{-1}]$, with $\phi(0)=1$ and such that, for all integer j, $\|\phi^{(j)}\|_\infty\ll_j\Delta^j$. Then, we introduce

$$W_h(x,y)=g_h(x,y)\phi(\ell_1 x-\ell_2 y-h).$$

In addition, we introduce another smooth function $w:\mathbb{R}\to\mathbb{R}$ compactly supported in $[Q,2Q]$ and such that $\|w^{(j)}\|_\infty\ll_j Q^{-j}$. If φ denotes Euler's phi function, let

$$\Lambda=\sum_{q\equiv 0[N\ell_1\ell_2]}w(q)\varphi(q)\asymp\frac{Q^2}{N\ell_1\ell_2}.$$

As a result, we can rewrite

$$\Sigma(\ell_1,\ell_2,H,a)=\sum_{h=1}^{H}a(h)\int_0^1\sum_{m_1,m_2\geq 1}\lambda_f(m_1)\overline{\lambda_f(m_2)}e(\ell_1 m_1\alpha)e(-\ell_2 m_2\alpha)W_h(m_1,m_2)e(-h\alpha)d\alpha.$$

By the Jutila's circle method (see [Blo05], lemma 3.1), we build an approximation $\tilde{I}$ to the characteristic function on $[0,1]$, which splits the α-integral in two parts, according to whether α is in a minor arc or a major arc. We easily estimate the contribution of minor arcs using our bound for the $L^2([0,1])$-norm of $1-\tilde{I}$ and we can write the contribution of major arcs of the shape

$$\frac{1}{2\delta\Lambda}\sum_{h=1}^{H}a(h)\sum_{q\equiv 0[N\ell_1\ell_2]}w(q)\sideset{}{^*}\sum_{d(q)}\int_{-\delta}^{\delta}\sum_{m_1,m_2\geq 1}\lambda_f(m_1)\overline{\lambda_f(m_2)}$$
$$e\left(\ell_1 m_1\left(\frac{d}{q}+\eta\right)\right)e\left(-\ell_2 m_2\left(\frac{d}{q}+\eta\right)\right)W_h(m_1,m_2)e\left(-h\left(\frac{d}{q}+\eta\right)\right)d\eta.$$

We transform short sums of exponentials in long sums of exponentials by means of a Voronoï summation formula (see [Blo05], lemma 2.2 or [KMV02], appendix A). If $S(m,n;q)$ denotes the classical Kloosterman sum, the major arcs contribution becomes

$$\frac{1}{2\delta\Lambda}\sum_{h=1}^{H}a(h)\sum_{q\equiv 0[N\ell_1\ell_2]}w(q)\int_{-\delta}^{\delta}e(-\eta h)$$
$$\times\sum_{m_1,m_2\geq 1}\lambda_f(m_1)\overline{\lambda_f(m_2)}S(-h,\ell_2 m_2-\ell_1 m_1;q)G_{q,h,\eta}(m_1,m_2)d\eta$$

where

$$G_{q,h,\eta}(x_1,x_2)=\frac{4\pi^2\ell_1\ell_2}{q^2}\int_0^{+\infty}\int_0^{+\infty}W_h(t_1,t_2)e(\ell_1t_1\eta-\ell_2t_2\eta)\\ \times J_{k-1}\left(\frac{4\pi\ell_1\sqrt{x_1t_1}}{q}\right)J_{k-1}\left(\frac{4\pi\ell_2\sqrt{x_2t_2}}{q}\right)dt_1dt_2.$$

We split this sum according to the value of $\ell_2m_2-\ell_1m_1$. It comes a diagonal contribution when $\ell_1m_1=\ell_2m_2$, which is easily bounded, and an off-diagonal contribution when $\ell_2m_2\neq\ell_1m_1$. It remains to estimate this off-diagonal contribution, which can be rewritten

$$\frac{\pi\ell_1\ell_2}{2\Lambda}\int_{-\delta}^{\delta}\int_0^{+\infty}\int_0^{+\infty}\sum_{h=1}^{H}a_{\eta,t_1,t_2}(h)\sum_{r\neq0}\sum_{\substack{1\leq m_1\leq\mathcal{M}_1\\1\leq m_2\leq\mathcal{M}_2\\ \ell_2m_2-\ell_1m_1=r}}\lambda_f(m_1)\overline{\lambda_f(m_2)}\\ \times\sum_{q\equiv0[N\ell_1\ell_2]}\frac{S(-h,r;q)}{q}\Phi_{t_1,t_2}\left(\frac{4\pi\sqrt{h|r|}}{q};m_1,m_2,r,h\right)dt_1dt_2d\eta.$$

where

$$a_{\eta,t_1,t_2}(h)=a(h)e(-\eta h)W_h(t_1,t_2)e(\ell_1t_1\eta-\ell_2t_2\eta)$$

$$\text{and }\Phi_{t_1,t_2}(x;m_1,m_2,r,h)=\frac{Qx}{\sqrt{h|r|}}J_{k-1}\left(\frac{x\ell_1\sqrt{m_1t_1}}{\sqrt{h|r|}}\right)J_{k-1}\left(\frac{x\ell_2\sqrt{m_2t_2}}{\sqrt{h|r|}}\right)w\left(\frac{4\pi\sqrt{h|r|}}{x}\right)w_1(h)$$

and with

$$\mathcal{M}_1=\frac{Q^2P_1^2k^2}{\ell_1^2M_1}(\ell_1\ell_2M_1M_2P_1P_2)^{\varepsilon}\text{ and }\mathcal{M}_2=\frac{Q^2P_2^2k^2}{\ell_2^2M_2}(\ell_1\ell_2M_1M_2P_1P_2)^{\varepsilon}.$$

The asymptotic behaviour of $G_{q,h,\eta}$ allows us to restrict sums over m_1 and m_2 to sums over $m_1\leq\mathcal{M}_1$ and $m_2\leq\mathcal{M}_2$. Applying the Kuznetsov trace formula (see [Blo05], lemma 2.4 or [DI83], theorem 1), we decompose this off-diagonal term as a sum of three terms: the contribution of discrete spectrum, the contribution of continuous spectrum and the contribution of holomorphic cusp forms. All of them will be evaluated by means of large sieve inequalities (see [Blo05], lemma 2.5 or [DI83], theorem 2). In the discrete spectrum, it may have exceptional eigenvalues even thought the Selberg's conjecture predicts that such exceptional eigenvalues do not exist. We prove that the contribution of non-exceptional eigenvalues, called real spectrum, is bounded by

$$A^{1/2}\|a\|_2h_1^{\theta}(P_1+P_2)^{\frac{3}{2}}\left(\sqrt{P_1+P_2}+\sqrt{\frac{H(h_1,\ell_1\ell_2)}{h_1\ell_1\ell_2}}\right)Q^{\varepsilon}$$

whereas we bound the contribution of exceptional eigenvalues by

$$A^{\frac{1}{2}+\theta}h_1^{\theta}\|a\|_2(P_1+P_2)^{\frac{3}{2}-\theta}\left(1+\sqrt{\frac{H(h_1,\ell_1\ell_2)}{h_1\ell_1\ell_2}}\right)Q^{\varepsilon}.$$

In addition, we show that the contribution of the continuous spectrum and of holomorphic cusp forms are bounded by

$$A^{1/2}\|a\|_2(P_1+P_2)^{3/2}\left(\sqrt{P_1+P_2}+\sqrt{\frac{H(h_1,\ell_1\ell_2)}{h_1\ell_1\ell_2}}\right)Q^{\varepsilon},$$

which is smaller than the estimate of the real spectrum. The result comes easily from these last three estimates.

□

Remark 4 In [Blo05], a factor $\left(\frac{LM}{HP}\right)^{\theta}$ appears, which could become $\left(\frac{A}{H(P_1+P_2)}\right)^{\theta}$ in our theorem, and which comes from the contribution of possibly exceptional eigenvalues in the discrete spectrum. However, we only find $\left(\frac{A}{P_1+P_2}\right)^{\theta}$.

Let us determine the required bounds for the test function in our case.

Lemma 5 *Let $\alpha,\beta \ll L^{-1}$ be complex numbers and let σ be any positive real number. For all non negative integers i and j, we have*

$$x^i y^j \frac{\partial^{i+j}F_{h;\ell_1,\ell_2}}{\partial x^i\partial y^j}(x,y) \ll_{i,j} \left(\frac{a}{A_{\ell_1}}\right)^{\frac{1}{2}+\Re\alpha+\sigma}\left(\frac{b}{A_{\ell_2}}\right)^{\frac{1}{2}+\Re\beta+\sigma} T^{1+2\sigma}(\ln T)^j$$

and the implicit constant does not depend on h.

Proof: Let $P_0(Y)=1$ and $P_j(Y)=\prod_{\ell=0}^{j-1}(Y-\ell)$ for $j\geq 1$. Let $\Re s=\sigma>0$, then we have

$$\begin{aligned} y^j\frac{\partial^j}{\partial y^j}\left(\int_{\mathbb{R}} w(t)\left(1+\frac{h}{y}\right)^{-it} g_{\alpha,\beta}(s,t)dt\right) &= \int_{\mathbb{R}} w(t)g_{\alpha,\beta}(s,t)\left(1+\frac{h}{y}\right)^{-it}Q_j(t)dt \\ &= \int_{\mathbb{R}} \frac{\left(1+\frac{h}{y}\right)^{-it}}{\left[i\ln\left(1+\frac{h}{y}\right)\right]^j}\frac{\partial^j}{\partial t^j}\left[w(t)g_{\alpha,\beta}(s,t)Q_j(t)\right]dt \end{aligned}$$

where $Q_j(t)=\sum_{r=0}^{j}\binom{j}{r}P_r(it)P_{j-r}(-it)\left(1+\frac{h}{y}\right)^{-(j-r)}$. Since $Q_j^{(r)}(t)\ll\left|\frac{h/y}{1+h/y}\right|^j t^{j-r}$ for $r\leq j$ and $\frac{\partial^r}{\partial t^r}\left[w(t)g_{\alpha,\beta}(s,t)\right]dt \ll T^{2\sigma}\Delta^{-r}$, we get

$$\frac{\partial^j}{\partial t^j}\left[w(t)g_{\alpha,\beta}(s,t)Q_j(t)\right] \ll_{s,j} T^{2\sigma}\left|\frac{h/y}{1+h/y}\right|^j(\ln T)^j.$$

and the implicit constant depends polynomially on s. Since $h/y\ll T^{-\gamma}$ in our range of summation, then

$$y^j\frac{\partial^j}{\partial y^j}\left(\int_{\mathbb{R}} w(t)\left(1+\frac{h}{y}\right)^{-it}g_{\alpha,\beta}(s,t)dt\right) \ll_{s,j} T^{1+2\sigma}(\ln T)^j \tag{18}$$

and the implicit constant does not depend on h. Writing

$$F_{h;\ell_1,\ell_2}(x,y)=\frac{\rho_{\ell_1}(x)\rho_{\ell_2}(y)}{2i\pi}\int_{(\sigma)}\frac{G(s)}{s}\left(\frac{a}{x}\right)^{\frac{1}{2}+\alpha+s}\left(\frac{b}{y}\right)^{\frac{1}{2}+\beta+s}\int_{\mathbb{R}}w(t)\left(1+\frac{h}{y}\right)^{-it}g_{\alpha,\beta}(s,t)dtds,$$

the result comes easily from the Leibniz formula and the bound (18).

□

Remark 5 The trivial bound for shifted convolution sums, namely taking absolute values and applying the Ramanujuan-Petersson bound on average, is given, for all $\varepsilon>0$, by

$$\sum_{\ell_1 m_1-\ell_2 m_2=h}\lambda_f(m_1)\lambda_f(m_2)g_h(m_1,m_2)\ll_\varepsilon \min\{M_1,M_2\}(M_1M_2)^\varepsilon. \tag{19}$$

The trivial bound (19) and lemma 5 imply the following corollary.

Corollary 4 *For all $\varepsilon>0$, we have*

$$I_{a,b}^{ND_1}(\alpha,\beta)\ll_\varepsilon \min\{a,b\}T^{1+\varepsilon}.$$

Remark 6 This trivial bound of $I_{a,b}^{ND_1}(\alpha,\beta)$ fails to prove Proposition 1. In other words, taking care of the oscillations of the Hecke eigenvalues is required to proving Proposition 1. In a first time, we apply the following bound, proved by Blomer in [Blo04], about shifted convolution sums.

Theorem 4 (Blomer, [Blo04]) *Let $\varepsilon>0$, and let ℓ_1, ℓ_2, h be positive integers. Let M_1, M_2, P_1, P_2 be real numbers greater than 1. Let g_h be a smooth function, supported on $[M_1,2M_1]\times[M_2,2M_2]$ such that $||g_h^{(ij)}||_\infty\ll_{i,j}(P_1/M_1)^i(P_2/M_2)^j$ for all $i,j\geq 0$. Then*

$$\sum_{\ell_1 m_1-\ell_2 m_2=h}\lambda_f(m_1)\lambda_f(m_2)g_h(m_1,m_2)\ll_{\varepsilon,P_1,P_2,N,k}(\ell_1M_1+\ell_2M_2)^{1/2+\theta+\varepsilon}.$$

This bound is uniform in ℓ_1, ℓ_2, h and the dependence on P_1, P_2, N and k is polynomial.

Remark 7 Remembering the trivial bound (19), this theorem agrees with the square-root cancellation philosophy.

As a consequence, theorem 4 and lemma 5 imply the following proposition, which gives a first admissible bound, and which will be improved in proposition 3.

Proposition 2 *Let $\alpha,\beta\ll L^{-1}$ be complex numbers and a,b be positive integers. For all $\varepsilon>0$, we have*

$$I_{a,b}^{ND_1}(\alpha,\beta)\ll_\varepsilon (ab)^{\frac{3}{4}+\frac{\theta}{2}}T^{\frac{1}{2}+\theta+\varepsilon}.$$

Proof: By theorem 4, lemma 5 gives

$$\sum_{am-bn=h}\lambda_f(m)\lambda_f(n)F_{h;\ell_1,\ell_2}(am,bn)\ll\left(\frac{a}{A_{\ell_1}}\right)^{\frac{1}{2}+\Re\alpha+\sigma}\left(\frac{b}{A_{\ell_2}}\right)^{\frac{1}{2}+\Re\beta+\sigma}T^{1+2\sigma}(\ln T)^\kappa(A_{\ell_1}+A_{\ell_2})^{\frac{1}{2}+\theta+\varepsilon}$$

where κ is a constant. Thus, thanks to lemma 4 and with $\frac{1}{2}+\theta+\varepsilon-\Re\alpha-\Re\beta-2\sigma>0$, we get

$$\begin{aligned}
I_{a,b}^{ND_1}(\alpha,\beta) &\ll T^{1-\gamma+2\sigma}(\ln T)^{\kappa} \sum_{\substack{A_{\ell_1}A_{\ell_2}\ll abT^{2+\varepsilon}\\ A_{\ell_1}\asymp A_{\ell_2}\\ A_{\ell_1},A_{\ell_2}\gg T^{\gamma}}} \left(\frac{a}{A_{\ell_1}}\right)^{\frac{1}{2}+\Re\alpha+\sigma}\left(\frac{b}{A_{\ell_2}}\right)^{\frac{1}{2}+\Re\beta+\sigma}\sqrt{A_{\ell_1}A_{\ell_2}}(A_{\ell_1}+A_{\ell_2})^{1/2+\theta+\varepsilon}\\
&\ll T^{1-\gamma+2\sigma}(ab)^{1/2+\sigma}(\ln T)^{\kappa}\sum_{T^{\gamma}\ll A_{\ell_1}\ll\sqrt{ab}T^{1+\varepsilon/2}} A_{\ell_1}^{\frac{1}{2}+\theta+\varepsilon-\Re\alpha-\Re\beta-2\sigma}\\
&\ll T^{1-\gamma}T^{\frac{1}{2}+\theta+\varepsilon}(ab)^{\frac{3}{4}+\frac{\theta}{2}}\sum_{T^{\gamma}\ll A_{\ell_1}\ll\sqrt{ab}T^{1+\varepsilon/2}} 1.
\end{aligned}$$

Then, we obtain $\sum_{T^{\gamma}\ll A_{\ell_1}\ll\sqrt{ab}T^{1+\varepsilon/2}}1=\sum_{1\leq 2^{\ell_1/2}\ll\sqrt{ab}T^{1-\gamma+\varepsilon/2}}1\ll\ln T$. Finally, the result easily comes from the choice $\gamma=1-\varepsilon$.

□

Remark 8 We are tempted to solve the shifted convolution problem on average (over h) and to take care of the resulting additional oscillations of the Hecke eigenvalues. For instance, using Theorem 6.3 of [Ric06], one can check that, for all $\varepsilon>0$,

$$I_{a,b}^{ND_1}(\alpha,\beta)\ll_{\varepsilon}(ab)^{\frac{3}{4}+\frac{\theta}{2}}T^{\frac{3}{2}+\theta+\varepsilon}.$$

It turns out that this bound is not admissible. It can be explained by the fact that the length of the h-sum is very small. That is why we need a bound for short sums of shifted convolution sums, which is given by theorem 3.

Proposition 3 *Let $\alpha,\beta\ll L^{-1}$ be complex numbers and a,b be positive integers. For all $\varepsilon>0$, we have*

$$I_{a,b}^{ND_1}(\alpha,\beta)\ll_{\varepsilon}(ab)^{\frac{1+\theta}{2}}T^{\frac{1}{2}+\theta+\varepsilon}.$$

Proof: We apply theorem 3 with $H=T^{-\gamma}\sqrt{A_{\ell_1}A_{\ell_2}}$, $h_1=1$ and $a(h)=\begin{cases}1 \text{ if } h\leq H,\\ 0 \text{ otherwise.}\end{cases}$

Therefore, from lemma 4, we get

$$\begin{aligned}
&I_{a,b}^{ND_1}(\alpha,\beta)\\
&\ll \sum_{\substack{A_{\ell_1}A_{\ell_2}\ll abT^{2+\varepsilon}\\ A_{\ell_1}\asymp A_{\ell_2}\\ A_{\ell_1},A_{\ell_2}\gg T^{\gamma}}} \left(\frac{a}{A_{\ell_1}}\right)^{\frac{1}{2}+\Re\alpha+\sigma}\left(\frac{b}{A_{\ell_2}}\right)^{\frac{1}{2}+\Re\beta+\sigma}T^{1+2\sigma+\varepsilon}\sqrt{A_{\ell_1}H}\left[\sqrt{\ln T}+\left(\frac{A_{\ell_1}}{\ln T}\right)^{\theta}\left(1+\sqrt{\frac{H}{ab}}\right)\right]\\
&\ll (ab)^{\frac{1}{2}+\theta}T^{1+2\sigma+\varepsilon-\frac{\gamma}{2}}\sum_{T^{\gamma}\ll A_{\ell}\ll\sqrt{ab}T^{1+\varepsilon}} A_{\ell}^{\theta-(\Re\alpha+\Re\beta+2\sigma)}\ll(ab)^{\frac{1+\theta}{2}}T^{1-\frac{\gamma}{2}+\theta+\varepsilon}.
\end{aligned}$$

Finally, the result comes from $\gamma=1-\varepsilon$.

□

Corollary 5 *Let $\alpha,\beta\ll L^{-1}$ be complex numbers and a,b be positive integers. For all $\varepsilon>0$, we have*

$$I_{a,b}^{ND_2}(\alpha,\beta)\ll_{\varepsilon}(ab)^{\frac{1+\theta}{2}}T^{\frac{1}{2}+\theta+\varepsilon}.$$

Proof: Set

$$w_1(t) = w(t)\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}\left(1+\frac{i(\alpha^2-\beta^2)}{t}\right).$$

Thanks to relation (9), we may write

$$I_{a,b}^{ND_2}(\alpha,\beta) = \sum_{am\neq bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}-\beta}n^{\frac{1}{2}-\alpha}}\int_{-\infty}^{+\infty} w_1(t)\left(\frac{am}{bn}\right)^{-it} V_{-\beta,-\alpha}(mn,t)dt + O\left(\frac{1}{T}\sum_{mn\ll T^{2+\varepsilon}}\frac{|\lambda_f(m)\lambda_f(n)|}{m^{\frac{1}{2}-\Re\beta}n^{\frac{1}{2}-\Re\alpha}}\right).$$

The error term becomes $O(T^{\varepsilon})$ and since w_1 satisfies (8a), (8b) and (8c) we may apply proposition 3 up to changing w by w_1 and (α,β) by $(-\beta,-\alpha)$.

□

3.1.3 Proof of proposition 1

Using proposition 3 and corollary 5, we trivially bound the off-diagonal term by

$$\begin{aligned}\sum_{a,b\leq M}\frac{\mu_f(a)\mu_f(b)}{\sqrt{ab}}P\left(\frac{\ln M/a}{\ln M}\right)P\left(\frac{\ln M/b}{\ln M}\right)\left[I_{a,b}^{ND_1}(\alpha,\beta)+I_{a,b}^{ND_2}(\alpha,\beta)\right] &\ll T^{\frac{1}{2}+\theta+\varepsilon}\sum_{a,b\leq M}(ab)^{\frac{\theta}{2}}\\ &\ll T^{\frac{1}{2}+\theta+\varepsilon}T^{\nu(2+\theta)}.\end{aligned}$$

Thus, if $\nu < \frac{1-2\theta}{4+2\theta}$, the off-diagonal part of $I_f(\alpha,\beta)$ is bounded by $T^{1-\varepsilon}$.

3.2 Evaluation of the diagonal term

For $i=1$ or $i=2$, let

$$I_f^{D_i}(\alpha,\beta) = \sum_{a,b\leq M}\frac{\mu_f(a)\mu_f(b)}{\sqrt{ab}}P\left(\frac{\ln M/a}{\ln M}\right)P\left(\frac{\ln M/b}{\ln M}\right)I_{a,b}^{D_i}(\alpha,\beta). \tag{20}$$

We consider the diagonal part $I_f^D(\alpha,\beta)$ of the mollified second moment. Thus, we have

$$I_f^D(\alpha,\beta) = I_f^{D_1}(\alpha,\beta) + I_f^{D_2}(\alpha,\beta). \tag{21}$$

In this section, we prove the following proposition:

Proposition 4 *Let $0<\nu<1$. For complex numbers $\alpha,\beta \ll L^{-1}$ such that $|\alpha+\beta| \gg L^{-1}$, we have*

$$I_f^D(\alpha,\beta) = \widehat{w}(0)c(\alpha,\beta) + O(T(\ln L)^4/L)$$

where $c(\alpha,\beta)$ is defined in relation (1).

3.2.1 Initial lemmas

Lemma 6 *Set $\Omega_{\alpha,\beta}$ the subset of vectors (u,v,s) in $\mathbb{C}^3$ satisfying*

$$\left\{\begin{array}{l} \Re u+\Re v>-1/2,\\ \Re s>-\frac{1}{4}-\frac{\Re\alpha}{2}-\frac{\Re\beta}{2},\\ \Re u+\Re s>-\frac{1}{2}-\Re\alpha,\\ \Re v+\Re s>-\frac{1}{2}-\Re\beta. \end{array}\right.$$

We have

$$\sum_{\substack{a,b,m,n\geq 1\\ am=bn}}\frac{\mu_f(a)\mu_f(b)\lambda_f(m)\lambda_f(n)}{a^{\frac{1}{2}+v}b^{\frac{1}{2}+u}m^{\frac{1}{2}+\alpha+s}n^{\frac{1}{2}+\beta+s}}=\frac{L(f\times f,1+\alpha+\beta+2s)L(f\times f,1+u+v)}{L(f\times f,1+\alpha+u+s)L(f\times f,1+\beta+v+s)}A_{\alpha,\beta}(u,v,s)$$

where $A_{\alpha,\beta}(u,v,s)$ is given by an absolutely convergent Euler product on $\Omega_{\alpha,\beta}$.

Proof : In order to simplify, set

$$\mathcal{P}=\sum_{\substack{a,b,m,n\geq 1\\ am=bn}}\frac{\mu_f(a)\mu_f(b)\lambda_f(m)\lambda_f(n)}{a^{\frac{1}{2}+v}b^{\frac{1}{2}+u}m^{\frac{1}{2}+\alpha+s}n^{\frac{1}{2}+\beta+s}}.$$

Using relation (3), for any prime number p such that $p\nmid N$, we get,

$$\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+1})}{p^{\ell s}}=\lambda_f(p)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell s}}-\frac{1}{p^s}\sum_{\ell\geq 0}\frac{\lambda_f(p^{\ell+1})\lambda_f(p^\ell)}{p^{\ell s}}$$

and

$$\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+2})}{p^{\ell s}}=\lambda_f(p^2)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell s}}-\frac{\lambda_f(p)}{p^s}\sum_{\ell\geq 0}\frac{\lambda_f(p^{\ell+1})\lambda_f(p^\ell)}{p^{\ell s}}.$$

Thus, we deduce

$$\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+1})}{p^{\ell s}} = \lambda_f(p)\left(1+\frac{1}{p^s}\right)^{-1}\sum_{l\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell s}} \quad (22)$$

$$\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+2})}{p^{\ell s}} = \left(1+\frac{1}{p^s}\right)^{-1}\left(\lambda_f(p^2)-\frac{1}{p^s}\right)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell s}}. \quad (23)$$

In addition, since we write

$$\mathcal{P}=\prod_p\left(\sum_{\substack{\ell_1,\ell_2,\ell_3,\ell_4\geq 0\\ \ell_1+\ell_3=\ell_2+\ell_4}}\frac{\mu_f(p^{\ell_1})\mu_f(p^{\ell_2})\lambda_f(p^{\ell_3})\lambda_f(p^{\ell_4})}{p^{\ell_1(\frac{1}{2}+v)}p^{\ell_2(\frac{1}{2}+u)}p^{\ell_3(\frac{1}{2}+\alpha+s)}p^{\ell_4(\frac{1}{2}+\beta+s)}}\right)$$

using relations (4) and (5), we get

$$\begin{aligned}\mathcal{P} = \prod_{p\nmid N}\Bigg[&\left(1+\frac{\lambda_f(p)^2}{p^{1+u+v}}+\frac{1}{p^{2(1+u+v)}}\right)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell(1+\alpha+\beta+2s)}}\\ &-\lambda_f(p)\left(\frac{1}{p^{1+v+\beta+s}}+\frac{1}{p^{1+u+\alpha+s}}\right)\left(1+\frac{1}{p^{1+u+v}}\right)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+1})}{p^{\ell(1+\alpha+\beta+2s)}}\\ &+\left(\frac{1}{p^{2(1+v+\beta+s)}}+\frac{1}{p^{2(1+u+\alpha+s)}}\right)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)\lambda_f(p^{\ell+2})}{p^{\ell(1+\alpha+\beta+2s)}}\Bigg]\\ &\times\prod_{p\mid N}\left[\left(1+\frac{\lambda_f(p)^2}{p^{1+u+v}}-\frac{\lambda_f(p)^2}{p^{1+\alpha+u+s}}-\frac{\lambda_f(p)^2}{p^{1+\beta+v+s}}\right)\sum_{\ell\geq 0}\frac{\lambda_f(p^\ell)^2}{p^{\ell(1+\alpha+\beta+2s)}}\right].\end{aligned}$$

Thus, it comes from relations (22) and (23),

$$\begin{aligned}\mathcal{P} &= L(f\times f,1+\alpha+\beta+2s)\prod_{p\mid N}\left[1+\frac{\lambda_f(p)^2}{p^{1+u+v}}-\frac{\lambda_f(p)^2}{p^{1+\alpha+u+s}}-\frac{\lambda_f(p)^2}{p^{1+\beta+v+s}}\right]\\ &\quad\times\prod_{p\nmid N}\Bigg[\left(1+\frac{\lambda_f(p)^2}{p^{1+u+v}}+\frac{1}{p^{2(1+u+v)}}\right)\left(1-\frac{1}{p^{2(1+\alpha+\beta+2s)}}\right)\\ &\quad-\lambda_f(p)^2\left(1-\frac{1}{p^{1+\alpha+\beta+2s}}\right)\left(\frac{1}{p^{1+v+\beta+s}}+\frac{1}{p^{1+u+\alpha+s}}\right)\left(1+\frac{1}{p^{1+u+v}}\right)\\ &\quad+\left(1-\frac{1}{p^{1+\alpha+\beta+2s}}\right)\left(\lambda_f(p^2)-\frac{1}{p^{1+\alpha+\beta+2s}}\right)\left(\frac{1}{p^{2(1+v+\beta+s)}}+\frac{1}{p^{2(1+u+\alpha+s)}}\right)\Bigg]\\ &= L(f\times f,1+\alpha+\beta+2s)\prod_{p}\left[1+\frac{\lambda_f(p)^2}{p^{1+u+v}}-\frac{\lambda_f(p)^2}{p^{1+\alpha+u+s}}-\frac{\lambda_f(p)^2}{p^{1+\beta+v+s}}+\chi_0(p)E_p\right]\end{aligned}$$

where

$$\begin{aligned}E_p = &\\ &\frac{1}{p^2}\left[\frac{1}{p^{2(u+v)}}-\frac{1}{p^{2(\alpha+\beta+2s)}}-\frac{\lambda_f(p)^2}{p^s}\left(\frac{1}{p^{u+\alpha}}+\frac{1}{p^{v+\beta}}\right)\left(\frac{1}{p^{u+v}}-\frac{1}{p^{\alpha+\beta+2s}}\right)+\frac{\lambda_f(p^2)}{p^{2s}}\left(\frac{1}{p^{2(u+\alpha)}}+\frac{1}{p^{2(v+\beta)}}\right)\right]\\ &+\frac{\lambda_f(p)^2}{p^{3+\alpha+\beta+3s}}\left[\frac{1}{p^{u+v}}\left(\frac{1}{p^{u+\alpha}}+\frac{1}{p^{v+\beta}}\right)-\frac{1}{p^s}\left(\frac{1}{p^{2(u+\alpha)}}+\frac{1}{p^{2(v+\beta)}}\right)-\frac{1}{p^{u+v+\alpha+\beta+s}}\right]\\ &+\frac{1}{p^{4+2(\alpha+\beta+2s)}}\left[\frac{1}{p^{2(u+\alpha+s)}}+\frac{1}{p^{2(v+\beta+s)}}-\frac{1}{p^{2(u+v)}}\right].\end{aligned}$$

Since the Rankin-Selberg L-function $L(f\times f,z)$ admits, for $\Re z>1$, the Euler product

$$L(f\times f,z)=\prod_{p\in\mathcal{P}}L_p(f\times f,z)$$

with

$$\begin{aligned}L_p(f\times f,z) &= \left(1-\frac{\alpha_f(p)^2}{p^z}\right)^{-1}\left(1-\frac{\alpha_f(p)\beta_f(p)}{p^z}\right)^{-2}\left(1-\frac{\beta_f(p)^2}{p^z}\right)^{-1}\\ &= \left(1-\frac{\lambda_f(p)^2}{p^z}+\chi_0(p)\left[\frac{2+\lambda_f(p)^2}{p^{2z}}-\frac{\lambda_f(p)^2}{p^{3z}}+\frac{1}{p^{4z}}\right]\right)^{-1},\end{aligned}$$

we may write

$$1+\frac{\lambda_f(p)^2}{p^{1+u+v}}-\frac{\lambda_f(p)^2}{p^{1+\alpha+u+s}}-\frac{\lambda_f(p)^2}{p^{1+\beta+v+s}}+\chi_0(p)E_p=\frac{L_p(f\times f,1+u+v)}{L_p(f\times f,1+u+\alpha+s)L_p(f\times f,1+v+\beta+s)}$$
$$\times\left[1+L_p(f\times f,1+u+\alpha+s)L_p(f\times f,1+v+\beta+s)\sum_{r=2}^{8}\sum_{\ell}\frac{a_{r,\ell}(p)}{p^{r+X_{r,\ell}(u,v,\alpha,\beta,s)}}\right]$$

where the sum over ℓ is finite, $X_{r,\ell}$ are linear forms in u,v,α,β,s and $a_{r,\ell}(p)$ are complex numbers with $|a_{r,\ell}(p)|\ll 1$. As a result, we obtain

$$\mathcal{P}=\frac{L(f\times f,1+\alpha+\beta+2s)L(f\times f,1+u+v)}{L(f\times f,1+\alpha+u+s)L(f\times f,1+\beta+v+s)}A_{\alpha,\beta}(u,v,s)$$

where

$$A_{\alpha,\beta}(u,v,s)=\prod_{p\in\mathcal{P}}\left[1+\sum_{r,\ell}O\left(\frac{1}{p^{r+X_{r,\ell}(\Re u,\Re v,\Re\alpha,\Re\beta,\Re s)}}\right)\right].$$

Then $A_{\alpha,\beta}(u,v,s)$ is an absolutely convergent Euler product in $\{\Re(\alpha+u+s)>-1\}\cap\{\Re(\beta+v+s)>-1\}\cap_{r,\ell}\{X_{r,\ell}(\Re u,\Re v,\Re\alpha,\Re\beta,\Re s)>1-r\}$. Making explicit all the linear forms $X_{2,\ell}$, we obtain $\{X_{2,\ell}(\Re u,\Re v,\Re\alpha,\Re\beta,\Re s)>-1\}=\Omega_{\alpha,\beta}$. Similarly, writing all the linear forms $X_{r,\ell}$, we prove that for (u,v,s) in $\Omega_{\alpha,\beta}$ then $X_{r,\ell}(\Re u,\Re v,\Re\alpha,\Re\beta,s)>-r/2\geq 1-r$. As a result, $A_{\alpha,\beta}(u,v,s)$ is an absolutely convergent Euler product on $\Omega_{\alpha,\beta}$ and defines an holomorphic function on $\Omega_{\alpha,\beta}$.

□

Lemma 7 *We have*

$$A_{0,0}(0,0,0)=1.$$

Proof: Thanks to lemma 6, if $\Re s>0$, we may write

$$A_{0,0}(s,s,s)=\sum_{\substack{a,b,m,n\geq 1\\ am=bn}}\frac{\mu_f(a)\mu_f(b)\lambda_f(m)\lambda_f(n)}{(ambn)^{\frac{1}{2}+s}}=\sum_{a,m\geq 1}\frac{\mu_f(a)\lambda_f(m)}{(am)^{1+2s}}\sum_{n|am}\mu_f(am/n)\lambda_f(n).$$

Since $(\mu_f(n))$ is the convolution inverse of $(\lambda_f(n))$, we have

$$\sum_{n|d}\mu_f(d/n)\lambda_f(n)=\delta(d).$$

Thus

$$A_{0,0}(s,s,s)=\mu_f(1)\lambda_f(1)=1.$$

To conclude, we extend this relation to $s=0$ by continuity in the half plane $\Re s>0$.

□

Lemma 8 *For all non negative integer a, we have*

$$\sum_{n\leq M}\frac{\lambda_f(n)^2}{n}\left(\ln\frac{M}{n}\right)^a=\frac{\underset{s=1}{Res}\,L(f\times f,s)}{\zeta^{(N)}(2)}\int_1^M\frac{1}{r}\left(\ln\frac{M}{r}\right)^a dr+O\left((\ln M)^a\right).$$

Proof: We may find in [Ran39] the following asymptotic behaviour

$$\sum_{n\leq x}\lambda_f(n)^2=x\frac{\underset{s=1}{Res}\,L(f\times f,s)}{\zeta^{(N)}(2)}+O(x^{3/5}).$$

That is why, after one integration by parts, we get

$$\sum_{n\leq M}\frac{\lambda_f(n)^2}{n}\left(\ln\frac{M}{n}\right)^a=\frac{\underset{s=1}{Res}\,L(f\times f,s)}{\zeta^{(N)}(2)}\sum_{n\leq M}\frac{1}{n}\left(\ln\frac{M}{n}\right)^a+O\left((\ln M)^a\right).$$

Finally, the Euler Maclaurin formula gives us

$$\sum_{n\leq M}\frac{1}{n}\left(\ln\frac{M}{n}\right)^a=\int_1^M\frac{1}{r}\left(\ln\frac{M}{r}\right)^a dr+O\left((\ln M)^a\right).$$

□

3.2.2 Estimation of $I_f^{D_1}(\alpha,\beta)$

For positive integers i and j, and for any positive real number δ, let

$$J_{\alpha,\beta}(i,j)=\frac{1}{(2i\pi)^2}\int_{(\delta)}\int_{(\delta)}M^{u+v}\frac{L(f\times f,1+u+v)A_{\alpha,\beta}(u,v,0)}{L(f\times f,1+\alpha+u)L(f\times f,1+\beta+v)}\frac{du}{u^{i+1}}\frac{dv}{v^{j+1}}. \tag{24}$$

Lemma 9 *Let $\alpha,\beta\ll 1/L$ be complex numbers. For positive integers i and j, we have*

$$J_{\alpha,\beta}(i,j)=\frac{(\ln M)^{i+j-1}}{i!j!\underset{s=1}{Res}\,L(f\times f,s)}\frac{d^2}{dxdy}\left[M^{\alpha x+\beta y}\int_0^1(x+u)^i(y+u)^j du\right]\Bigg|_{x=y=0}$$
$$+O\left(L^{i+j-2}\left(1+\frac{(\ln L)^2}{L^{i-1}}\right)\left(1+\frac{(\ln L)^2}{L^{j-1}}\right)\right).$$

Proof: We use the Dirichlet series of $L(f\times f,s)$. Moving far to the right either u or v integration line, we obtain

$$J_{\alpha,\beta}(i,j)=\sum_{n\leq M}\frac{\lambda_f(n)^2}{n}\frac{1}{(2i\pi)^2}\int_{(\delta)}\int_{(\delta)}\left(\frac{M}{n}\right)^{u+v}\frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f,1+\alpha+u)L(f\times f,1+\beta+v)}\frac{du}{u^{i+1}}\frac{dv}{v^{j+1}}.$$

Let

$$r_{\alpha,\beta}^{(i,j)}(u,v)=\left(\frac{M}{n}\right)^{u+v}\frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f,1+\alpha+u)L(f\times f,1+\beta+v)}\frac{1}{u^{i+1}v^{j+1}}.$$

We consider the contour γ defined by $\gamma=\gamma_1\cup\gamma_2\cup\gamma_3$ with $c>0$, $Y\gg 1$ large and

$$\gamma_1=\{i\tau,\ |\tau|\geq Y\},\ \ \gamma_2=\left\{\sigma\pm iY,\ \frac{-c}{\ln Y}\leq\sigma\leq 0\right\}\ \text{and}\ \gamma_3=\left\{\frac{-c}{\ln Y}+i\tau,\ |\tau|\leq Y\right\}.$$

By the standard zero-free region of $L(f\times f,s)$, we move the integration lines $\Re u=\Re v=\delta$ to u and v on γ. Thus,

$$\begin{aligned}
&\frac{1}{(2i\pi)^2}\int_{(\delta)}\int_{(\delta)} r_{\alpha,\beta}^{(i,j)}(u,v)dudv\\
=&\operatorname*{Res}_{u=0}\frac{1}{2i\pi}\int_{\Re v=\delta} r_{\alpha,\beta}^{(i,j)}(u,v)dv+\frac{1}{(2i\pi)^2}\int_{u\in\gamma}\int_{\Re v=\delta} r_{\alpha,\beta}^{(i,j)}(u,v)dvdu\\
=&\operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v)+\operatorname*{Res}_{u=0}\frac{1}{2i\pi}\int_{v\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)dv+\operatorname*{Res}_{v=0}\frac{1}{2i\pi}\int_{u\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)du+\frac{1}{(2i\pi)^2}\int_\gamma\int_\gamma r_{\alpha,\beta}^{(i,j)}(u,v)dudv.
\end{aligned}$$

• We begin with the estimation of $\operatorname*{Res}_{u=0}\frac{1}{2i\pi}\int_{v\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)dv$. We express the residue as a contour integral on a circle with radius $1/L$. We get

$$\begin{aligned}
&\operatorname*{Res}_{u=0}\frac{1}{2i\pi}\int_{v\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)dv\\
=\ &\frac{1}{(2i\pi)^2}\int_{v\in\gamma}\frac{\left(\frac{M}{n}\right)^v}{L(f\times f,1+\beta+v)}\oint_{D(0,L^{-1})}\left(\frac{M}{n}\right)^u\frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f,1+\alpha+u)}\frac{du}{u^{i+1}}\frac{dv}{v^{j+1}}.
\end{aligned}$$

Furthermore, since $|\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)|\ll 1$ in our range of integration and since

$$\frac{1}{L(f\times f,1+\alpha+u)}\ll\alpha+u\ll L^{-1}\ \text{because}\ u\asymp 1/L,$$

we obtain

$$\operatorname*{Res}_{u=0}\frac{1}{2i\pi}\int_{v\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)dv\ll L^{i-1}\int_{v\in\gamma}\frac{\left(\frac{M}{n}\right)^{\Re v}}{|L(f\times f,1+\beta+v)|}\frac{dv}{|v|^{j+1}}.$$

As we have $-\frac{L'(f\times f,z)}{L(f\times f,z)}=\sum_{n\geq 1}\frac{\Lambda_f(n)}{n^z}$ with $\Lambda_f(n)\geq 0$, we deduce (confer [Ten95] part 3.10 and [IK04] part 5.3) that

$$\frac{1}{L(f\times f,\sigma+i\tau)}\ll\ln|\tau|,$$

it comes

$$\begin{aligned}
&\int_{v\in\gamma}\frac{\left(\frac{M}{n}\right)^{\Re v}}{|L(f\times f,1+\beta+v)|}\frac{dv}{|v|^{j+1}}\\
\ll\ &\int_{|\tau|\geq Y}\frac{\ln\tau}{|\tau|^{j+1}}d\tau+\ln Y\int_{-\frac{c}{\ln Y}}^0\frac{d\sigma}{|\sigma+iY|^{j+1}}+\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\ln Y\int_{|\tau|\leq Y}\frac{d\tau}{\left|\tau-i\frac{c}{\ln Y}\right|^{j+1}}\\
\ll\ &\frac{\ln Y}{Y^j}+(\ln Y)^{j+1}\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}.
\end{aligned}\tag{25}$$

As a result, we get the bound

$$\underset{u=0}{Res}\frac{1}{2i\pi}\int_{v\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)dv \ll L^{i-1}\ln Y\left(\frac{1}{Y^j}+(\ln Y)^j\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right). \tag{26}$$

• Since $r_{\alpha,\beta}^{(i,j)}(u,v)=r_{\beta,\alpha}^{(j,i)}(v,u)$, it comes immediately from the previous bound

$$\underset{v=0}{Res}\frac{1}{2i\pi}\int_{u\in\gamma} r_{\alpha,\beta}^{(i,j)}(u,v)du \ll L^{j-1}\ln Y\left(\frac{1}{Y^i}+(\ln Y)^i\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right). \tag{27}$$

• By the relation (25), we bound $\int_\gamma\int_\gamma r_{\alpha,\beta}^{(i,j)}(u,v)dudv$.

$$\begin{aligned}\frac{1}{(2i\pi)^2}\int_\gamma\int_\gamma r_{\alpha,\beta}^{(i,j)}(u,v)dudv &= \frac{1}{(2i\pi)^2}\int_\gamma\int_\gamma\left(\frac{M}{n}\right)^{u+v}\frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f,1+\alpha+u)L(f\times f,1+\beta+v)}\frac{1}{u^{i+1}v^{j+1}}\\
&\ll \int_\gamma\frac{\left(\frac{M}{n}\right)^{\Re v}}{|L(f\times f,1+\beta+v)|}\frac{dv}{|v|^{j+1}}\int_\gamma\frac{\left(\frac{M}{n}\right)^{\Re u}}{|L(f\times f,1+\alpha+u)|}\frac{du}{|u|^{i+1}}\\
&\ll \left(\frac{\ln Y}{Y^j}+(\ln Y)^{j+1}\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right)\left(\frac{\ln Y}{Y^i}+(\ln Y)^{i+1}\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right)\\
&\ll (\ln Y)^2\left(\frac{1}{Y^{i+j}}+(\ln Y)^{i+j}\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right).\end{aligned} \tag{28}$$

• In addition, for any positive integer ℓ, let

$$W_\ell=\sum_{n\le M}\frac{\lambda_f(n)^2}{n}\left(\frac{1}{Y^\ell}+(\ln Y)^\ell\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\right).$$

We can bound

$$\begin{aligned}W_\ell &\ll \frac{1}{Y^\ell}\sum_{n\le\frac{M}{(Y\ln Y)^{\ell(\ln Y)/c}}}\frac{\lambda_f(n)^2}{n}+(\ln Y)^\ell\sum_{\frac{M}{(Y\ln Y)^{\ell(\ln Y)/c}}\le n\le M}\frac{\lambda_f(n)^2}{n}\left(\frac{M}{n}\right)^{-\frac{c}{\ln Y}}\\
&\ll \frac{L}{Y^\ell}+(\ln Y)^\ell\sum_{0\le d\le\frac{\ln(Y\ln Y)}{\ln 2}}2^{-d\ell}\sum_{\frac{M}{(2^{d+1})^{\ell(\ln Y)/c}}\le n\le\frac{M}{(2^d)^{\ell(\ln Y)/c}}}\frac{\lambda_f(n)^2}{n}\\
&\ll \frac{L}{Y^\ell}+(\ln Y)^{\ell+1}.\end{aligned}$$

As a consequence, by relations (26), (27) and (28), we obtain

$$\begin{aligned}J_{\alpha,\beta}(i,j) &= \sum_{n\le M}\left[\frac{\lambda_f(n)^2}{n}\underset{u=v=0}{Res}\,r_{\alpha,\beta}^{(i,j)}(u,v)\right]+O\left(L^{i-1}W_j\ln Y+L^{j-1}W_i\ln Y+W_{i+j}(\ln Y)^2\right)\\
&= \sum_{n\le M}\left[\frac{\lambda_f(n)^2}{n}\underset{u=v=0}{Res}\,r_{\alpha,\beta}^{(i,j)}(u,v)\right]+O\left[L^{i-1}\ln Y\left(\frac{L}{Y^j}+(\ln Y)^j\right)+L^{j-1}\ln Y\left(\frac{L}{Y^i}+(\ln Y)^i\right)\right.\\
&\qquad\left.+(\ln Y)^2\left(\frac{L}{Y^{i+j}}+(\ln Y)^{i+j}\right)\right].\end{aligned}$$

Choosing $Y = L$, which is allowed, the error term becomes $O\left(\left(L^{i-1} + (\ln L)^2\right)\left(L^{j-1} + (\ln L)^2\right)\right)$. Then

$$J_{\alpha,\beta}(i,j) = \sum_{n \leq M} \left[\frac{\lambda_f(n)^2}{n} \operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v) \right] + O\left(L^{i+j-2} \left(1 + \frac{(\ln L)^2}{L^{i-1}}\right)\left(1 + \frac{(\ln L)^2}{L^{j-1}}\right)\right) \tag{29}$$

• We finish with the estimation of $\operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v)$. To do so again, we express the residue as a contour integral on a circle with radius $1/L$. Thus,

$$\operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v) = \frac{1}{(2i\pi)^2} \oint_{D(0,L^{-1})} \oint_{D(0,L^{-1})} \left(\frac{M}{n}\right)^{u+v} \frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f, 1+\alpha+u)L(f\times f, 1+\beta+v)} \frac{du}{u^{i+1}} \frac{dv}{v^{j+1}}$$

Furthermore, since $u \asymp v \asymp 1/L$, we have

$$\begin{aligned}
\zeta^{(N)}(2(1+u+v)) &= \zeta^{(N)}(2) + O(1/L), \\
A_{\alpha,\beta}(u,v,0) &= A_{0,0}(0,0,0) + O(1/L), \\
\frac{1}{L(f\times f, 1+\alpha+u)} &= \frac{\alpha+u}{\operatorname*{Res}_{s=1} L(f\times f,s)}(1+O(1/L)), \\
\frac{1}{L(f\times f, 1+\beta+v)} &= \frac{\beta+v}{\operatorname*{Res}_{s=1} L(f\times f,s)}(1+O(1/L)).
\end{aligned}$$

With lemma 7, we obtain

$$\frac{\zeta^{(N)}(2(1+u+v))A_{\alpha,\beta}(u,v,0)}{L(f\times f, 1+\alpha+u)L(f\times f, 1+\beta+v)} = (\alpha+u)(\beta+v)\frac{\zeta^{(N)}(2)}{[\operatorname*{Res}_{s=1} L(f\times f,s)]^2} + O(1/L^3).$$

Then

$$\operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v) = \frac{\zeta^{(N)}(2)}{[\operatorname*{Res}_{s=1} L(f\times f,s)]^2} \left[\frac{1}{2i\pi}\oint_{D(0,L^{-1})} \left(\frac{M}{n}\right)^u \frac{\alpha+u}{u^{i+1}}du\right]\left[\frac{1}{2i\pi}\oint_{D(0,L^{-1})} \left(\frac{M}{n}\right)^v \frac{\beta+v}{v^{j+1}}dv\right] + O\left(L^{i+j-3}\right).$$

In addition, thanks to Cauchy formula, we compute for any positive integer ℓ

$$\begin{aligned}
\frac{1}{2i\pi}\oint_{D(0,L^{-1})} \left(\frac{M}{n}\right)^u \frac{\alpha+u}{u^{\ell+1}}du &= \frac{d}{dx}\left[\frac{e^{\alpha x}}{2i\pi}\oint_{D(0,L^{-1})} \left(\frac{M}{n}e^x\right)^u \frac{du}{u^{\ell+1}}\right]\Bigg|_{x=0} \\
&= \frac{d}{dx}\left[e^{\alpha x}\frac{1}{\ell!}\frac{d^\ell}{du^\ell}\left[\left(\frac{M}{n}e^x\right)^u\right]\Bigg|_{u=0}\right]\Bigg|_{x=0} \\
&= \frac{1}{\ell!}\frac{d}{dx}\left[e^{\alpha x}\left(x+\ln\frac{M}{n}\right)^\ell\right]\Bigg|_{x=0} \\
&= \frac{1}{\ell!}\left[\alpha\left(\ln\frac{M}{n}\right)^\ell + \ell\left(\ln\frac{M}{n}\right)^{\ell-1}\right].
\end{aligned}$$

Then, we can write

$$\operatorname*{Res}_{u=v=0} r_{\alpha,\beta}^{(i,j)}(u,v) = \frac{\zeta^{(N)}(2)}{i!j![\operatorname*{Res}_{s=1} L(f\times f,s)]^2}\left[\alpha\beta\left(\ln\frac{M}{n}\right)^{i+j} + (\alpha i+\beta j)\left(\ln\frac{M}{n}\right)^{i+j-1} + ij\left(\ln\frac{M}{n}\right)^{i+j-2}\right] + O\left(L^{i+j-3}\right).$$

From (29) and by lemma 8, we get

$$J_{\alpha,\beta}(i,j)=\frac{[\underset{s=1}{Res}\,L(f\times f,s)]^{-1}}{i!j!}\int_1^M\left(\alpha\left(\ln\frac{M}{r}\right)^i+i\left(\ln\frac{M}{r}\right)^{i-1}\right)\left(\beta\left(\ln\frac{M}{r}\right)^j+j\left(\ln\frac{M}{r}\right)^{j-1}\right)\frac{dr}{r}$$
$$+O\left(L^{i+j-2}\left(1+\frac{(\ln L)^2}{L^{i-1}}\right)\left(1+\frac{(\ln L)^2}{L^{j-1}}\right)\right).$$

Changing the variable r to u with $r=M^{1-u}$ concludes the proof.

□

Lemma 10 *If $0<\nu<1$ and $\alpha,\beta\ll L^{-1}$ are complex numbers with $|\alpha+\beta|\gg L^{-1}$, then*

$$I_f^{D_1}(\alpha,\beta)=\frac{\widehat{w}(0)}{(\alpha+\beta)\ln M}\,\frac{d^2}{dxdy}\left[M^{\alpha x+\beta y}\int_0^1P(x+u)P(y+u)du\right]\Big|_{x=y=0}+O\left(\frac{T(\ln L)^4}{L}\right).$$

Proof: We use the Mellin transformation to write

$$\left(\frac{\ln M/a}{\ln M}\right)^i=\begin{cases}\frac{i!}{(\ln M)^i}\frac{1}{2i\pi}\int_{(1)}\left(\frac{M}{a}\right)^v\frac{dv}{v^{i+1}} & \text{if } 1\le a\le M,\\ 0 \quad \text{if } a>M.\end{cases}$$

Set $P(X)=\sum_{i=1}^{\deg P}a_iX^i$. Thus, from relation (20) which defines $I_f^{D_1}(\alpha,\beta)$, we get

$$I_f^{D_1}(\alpha,\beta)=\int_{-\infty}^{+\infty}w(t)\sum_{i,j\ge1}\frac{a_ia_ji!j!}{(\ln M)^{i+j}}\frac{1}{(2i\pi)^3}\int_{(1)}\int_{(1)}\int_{(1)}\frac{M^{u+v}G(s)}{s}g_{\alpha,\beta}(s,t)$$
$$\sum_{\substack{a,b,m,n\ge1\\ am=bn}}\frac{\mu_f(a)\mu_f(b)\lambda_f(m)\lambda_f(n)}{a^{\frac12+v}b^{\frac12+u}m^{\frac12+\alpha+s}n^{\frac12+\beta+s}}ds\frac{du}{u^{j+1}}\frac{dv}{v^{i+1}}dt.$$

Due to lemma 6, we can write

$$I_f^{D_1}(\alpha,\beta)=\int_{-\infty}^{+\infty}w(t)\sum_{i,j\ge1}\frac{a_ia_ji!j!}{(\ln M)^{i+j}}\frac{1}{(2i\pi)^3}\int_{(1)}\int_{(1)}\int_{(1)}\frac{M^{u+v}G(s)}{s}g_{\alpha,\beta}(s,t)A_{\alpha,\beta}(u,v,s)$$
$$\frac{L(f\times f,1+\alpha+\beta+2s)L(f\times f,1+u+v)}{L(f\times f,1+\alpha+u+s)L(f\times f,1+\beta+v+s)}ds\frac{du}{u^{j+1}}\frac{dv}{v^{i+1}}dt.$$

We specialize G in

$$G(s)=e^{s^2}\frac{(\alpha+\beta)^2-(2s)^2}{(\alpha+\beta)^2}.$$

As a result, $G(s)L(f\times f,1+\alpha+\beta+2s)$ is an entire function. First, we move the integration lines from $\Re u=$Re $\Re v=1$ to $\Re u=\Re v=\delta$, with δ small, in order to assure the absolute

convergence of $A_{\alpha,\beta}(u,v,s)$. Secondly, we move the integration line from $\Re s = 1$ to $\Re s = -\delta+\epsilon$, with $0 < \epsilon < \delta$, crossing a pole at $s = 0$. Since $t \asymp T$, $\nu < 1$ and $g_{\alpha,\beta}(s,t) \ll T^{2s}$, we can bound

$$\int_{-\infty}^{+\infty} w(t) \sum_{i,j} \frac{a_i a_j i! j!}{(\ln M)^{i+j}} \frac{1}{(2i\pi)^3} \int_{Re\, u=\delta} \int_{Re\, v=\delta} \int_{Re\, s=-\delta+\epsilon} \frac{M^{u+v} G(s)}{s} g_{\alpha,\beta}(s,t) A_{\alpha,\beta}(u,v,s)$$
$$\frac{L(f\times f, 1+\alpha+\beta+2s) L(f\times f, 1+u+v)}{L(f\times f, 1+\alpha+u+s) L(f\times f, 1+\beta+v+s)} ds \frac{du}{u^{j+1}} \frac{dv}{v^{i+1}} dt$$
$$\ll \int_{-\infty}^{+\infty} |w(t)| dt\, T^{2(-\delta+\epsilon)} M^{2\delta} \ll T^{1-(2-2\nu)\delta+\epsilon} \ll T^{1-\varepsilon}.$$

for sufficiently small ε. Then, using some previous notations, this estimation gives

$$I_f^{D_1}(\alpha,\beta) = \widehat{w}(0) L(f\times f, 1+\alpha+\beta) \sum_{i,j} \frac{a_i a_j i! j!}{(\ln M)^{i+j}} J_{\alpha,\beta}(i,j) + O\left(T^{1-\varepsilon}\right).$$

Thanks to lemma 9 and since

$$L(f\times f, 1+\alpha+\beta) = \frac{\underset{s=1}{Res}\, L(f\times f, s)}{\alpha+\beta} + O(1),$$

we get

$$I_f^{D_1}(\alpha,\beta) = \widehat{w}(0) \left[\frac{\underset{s=1}{Res}\, L(f\times f, s)}{\alpha+\beta} + O(1) \right]$$
$$\times \left[\frac{1}{\underset{s=1}{Res}\, L(f\times f, s) \ln M} \frac{d^2}{dxdy} \left[M^{\alpha x+\beta y} \int_0^1 P(x+u)P(y+u)du \right]\Big|_{x=y=0} + O\left(\frac{(\ln L)^4}{L^2} \right) \right] + O\left(T^{1-\varepsilon}\right)$$
$$= \frac{\widehat{w}(0)}{(\alpha+\beta)\ln M} \frac{d^2}{dxdy} \left[M^{\alpha x+\beta y} \int_0^1 P(x+u)P(y+u)du \right]\Big|_{x=y=0} + O\left(\frac{T(\ln L)^4}{(\alpha+\beta)L^2} \right) + O(T/L).$$

We conclude using the assumption $|\alpha+\beta| \gg L^{-1}$.

□

Remark 9 Sometimes, it may be useful to consider the relation

$$\frac{d^2}{dxdy} \left[M^{\alpha x+\beta y} \int_0^1 P(x+u)P(y+u)du \right]\Big|_{x=y=0} = \int_0^1 (P'(u)+\alpha \ln MP(u))(P'(u)+\beta \ln MP(u))du. \tag{30}$$

Lemma 11 *If $0 < \nu < 1$ and $\alpha, \beta \ll L^{-1}$ are complex numbers with $|\alpha+\beta| \gg L^{-1}$, then*

$$I_f^{D2}(\alpha,\beta) = T^{-2(\alpha+\beta)} I_f^{D1}(-\beta,-\alpha) + O(T/L).$$

Proof: We write

$$I_{a,b}^{D_2}(\alpha,\beta)=\sum_{am=bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac12-\beta}n^{\frac12-\alpha}}\int_{-\infty}^{+\infty}w(t)\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}V_{-\beta,-\alpha}(mn,t)dt+O\left(\sum_{\substack{am=bn\\ mn\ll T^{2+\varepsilon}}}\frac{|\lambda_f(m)\lambda_f(n)|}{m^{\frac12-\beta}n^{\frac12-\alpha}}\right).$$

Let $a'=a/(a,b)$ and $b'=b/(a,b)$. Then, for all $\delta>0$, the error term in the previous relation becomes

$$\sum_{\substack{am=bn\\ mn\ll T^{2+\varepsilon}}}\frac{|\lambda_f(m)\lambda_f(n)|}{m^{\frac12-\beta}n^{\frac12-\alpha}}\ll\frac{1}{(a'b')^{1/2-\delta}}\sum_{k\ll\frac{T^{1+\frac{\varepsilon}{2}}}{\sqrt{a'b'}}}\frac{1}{k^{1-\alpha-\beta-2\delta}}\ll\frac{T^{\varepsilon}}{\sqrt{a'b'}}$$

Therefore, if $w_2(t)=w(t)\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}$, we may write

$$I_f^{D_2}(\alpha,\beta)=\sum_{a,b\leq M}\frac{\mu_f(a)\mu_f(b)}{\sqrt{ab}}P\left(\frac{\ln M/a}{\ln M}\right)P\left(\frac{\ln M/b}{\ln M}\right)\left[\sum_{am=bn}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac12-\beta}n^{\frac12-\alpha}}\int_{-\infty}^{+\infty}w_2(t)V_{-\beta,-\alpha}(mn,t)dt\right]$$
$$+O\left(T^{\varepsilon}\sum_{a,b\leq M}\frac{1}{\sqrt{ab}\sqrt{a'b'}}\right).$$

We also have

$$\sum_{a,b\leq M}\frac{1}{\sqrt{ab}\sqrt{a'b'}}\ll\sum_{k\leq M}\frac1k\sum_{a,b\leq\frac{M}{k}}\frac{1}{ab}\ll\sum_{k\leq M}\frac1k\left(\ln\frac{M}{k}\right)^2\ll(\ln M)^3$$

and since w_2 satisfies (8a), (8b) and (8c), up to changing w by w_2 and (α,β) by $(-\beta,-\alpha)$, lemma 10 gives

$$I_f^{D_2}(\alpha,\beta)=\frac{\widehat{w_2}(0)}{(-\alpha-\beta)\ln M}\,\frac{d^2}{dxdy}\left[M^{-\alpha x-\beta y}\int_0^1P(x+u)P(y+u)du\right]\Big|_{x=y=0}+O\left(T(\ln L)^4/L\right).$$

To conclude, due to the support of w, we can write $\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}=T^{-2(\alpha+\beta)}+O(1/L)$ which gives $\widehat{w_2}(0)=T^{-2(\alpha+\beta)}\widehat{w}(0)+O(T/L)$.

□

3.2.3 Proof of proposition 4

From relation (21) and using lemma 11, we can write

$$\begin{aligned}I_f^D(\alpha,\beta)&=I_f^{D1}(\alpha,\beta)+T^{-2(\alpha+\beta)}I_f^{D1}(-\beta,-\alpha)+O(T(\ln L)^4/L)\\&=I_f^{D1}(\alpha,\beta)+I_f^{D1}(-\beta,-\alpha)+I_f^{D1}(-\beta,-\alpha)\left[T^{-2(\alpha+\beta)}-1\right]+O(T(\ln L)^4/L).\end{aligned}$$

Finally, using relation (30) and lemma 10, we have

$$I_f^{D1}(\alpha,\beta)+I_f^{D1}(-\beta,-\alpha)=\widehat{w}(0)+O(T(\ln L)^4/L).$$

Combining these relations, we get the result.

□

4 Effective proportion of zeros on the critical line

In this section, we prove corollary 1. From theorem 1, we may deduce the following theorem about mollified second moment of $L(f,s)$ and its derivative.

Theorem 5 *Let Q be a polynomial with complex coefficients satisfying $Q(0)=1$. Let*

$$V(s)=Q\left(-\frac{1}{2\ln T}\frac{d}{ds}\right)(L(f,s)).$$

Then, if $\nu<\frac{1-2\theta}{4+2\theta}$, we have

$$\frac{1}{T}\int_1^T|V\psi(\sigma_0+it)|^2dt=c(P,Q,2R,\nu/2)+o(1)$$

where

$$c(P,Q,r,\xi) \quad = \quad 1+\frac{1}{\xi}\int_0^1\int_0^1 e^{2rs}\left[\frac{d}{dx}\left(e^{r\xi x}Q(s+\xi x)P(x+u)\right)\Big|_{x=0}\right]^2 duds.$$

We do not give the proof of this theorem, which is essentially the same than the one of theorem 1 in [You10]. We may refer to part 2 and 3 of [You10] to find more details. Now, Q refers to a polynomial with complex coefficients of the shape

$$Q(x)=1+\sum_{n=1}^{M}i^{n+1}\lambda_n\left[(1-2x)^n-1\right] \tag{31}$$

where M is a positive integer and $(\lambda_1,...,\lambda_M)$ belongs to $\mathbb{R}^M$.

Let $N_f(T)$ (resp. $N_{f,0}(T)$) be the number of non-trivial zeros ρ (resp. on the critical line) of $L(f,s)$ with $0<\Im(\rho)\leq T$ for f a holomorphic primitive cusp form of even weight, square-free level and trivial character.

Proposition 5 *For f a holomorphic primitive cusp form of even weight, square-free level and trivial character, Q as in* (31)*, we have*

$$\liminf_{T\to+\infty}\frac{N_{f,0}(T)}{N_f(T)}\geq\limsup_{T\to+\infty}\left[1-\frac{1}{2R}\ln\left(\frac{1}{T}\int_1^T|V\psi(\sigma_0+it)|^2dt\right)\right]$$

Therefore, using theorem 5, we get

$$\liminf_{T\to+\infty}\frac{N_{f,0}(T)}{N_f(T)}\geq 1-\inf_{P,Q,R}\frac{1}{R}\ln c(P,Q,R,\nu/2).$$

The work of Kim and Sarnak ([Kim03]) gives $\theta=7/64$ then theorem 5 gives $\nu=5/27$. The Ramanujuan-Petersson conjecture ($\theta=0$) gives $\nu=1/4$.

Lemma 12 ([Con89], part 4) *We have*

$$\inf_P \frac{1}{R}\ln c(P,Q,R,\nu/2) = \frac{1}{R}\ln\left(\frac{1+|w(1)|^2}{2} + \frac{A\alpha}{\tanh\frac{\nu\alpha}{2}}\right)$$

where $w(x) = e^{Rx}Q(x)$, $A = \int_0^1 |w(x)|^2 dx$, $B = \int_0^1 w(x)\overline{w'(x)}dx$, $C = \int_0^1 |w'(x)|^2 dx$ *and* $\alpha = \frac{\sqrt{(B-\overline{B})^2+4AC}}{2A}$.

For empirical reasons, we restrict ourselves to polynomials Q with real coefficients of the shape

$$Q(x) = 1 + \sum_{n=1}^{N} h_n\left[(1-2x)^{2n-1} - 1\right] \tag{32}$$

where N is a positive integer and $(h_1, ..., h_N)$ belongs to $\mathbb{R}^N$. Then we get

$$\liminf_{T\to+\infty} \frac{N_{f,0}(T)}{N_f(T)} \geq 1 - \inf_{Q \text{ real},\ R} \frac{1}{R}\ln\left(\frac{1+w(1)^2}{2} + \frac{\sqrt{AC}}{\tanh\left(\frac{\nu}{2}\sqrt{\frac{C}{A}}\right)}\right). \tag{33}$$

To obtain corollary 1, we choose $N = 4$, R and Q as in (32) where R, h_1, h_2, h_3, h_4 are given in the following table.

	$\nu = \frac{1}{6}$	$\nu = \frac{5}{27}$	$\nu = \frac{1}{4}$
R	$6,6838894702116801322$	$6,4278834168344993342$	$5,6503610091685135131$
h_1	$1,6017785744634898860$	$1,5898336242677838745$	$1,5369390514358411982$
h_2	$-3,0362512753510924917$	-2.8999828229132398066	$-2,7929104872905007806$
h_3	$3,0757757634512927939$	3.0171733454035522056	$2,7758193765120241770$
h_4	$-1,1407980564855935531$	$-1,1164150244992046552$	$-1,0187870607687957034$

5 Non-mollified second integral moment

This section contains the proof of theorem 2. For more convenience, we set

$$M_{f,2}(\alpha,\beta) = \int_{-\infty}^{+\infty} w(t) L\left(f, \frac{1}{2}+\alpha+it\right) L\left(f, \frac{1}{2}+\beta-it\right) dt$$

where w satisfies (8a), (8b) and (8c). Applying the approximate equation (lemma 1) and using some previous notations, we can write

$$M_{f,2}(\alpha,\beta) = I_{1,1}^{D_1}(\alpha,\beta) + I_{1,1}^{D_2}(\alpha,\beta) + I_{1,1}^{ND_1}(\alpha,\beta) + I_{1,1}^{ND_2}(\alpha,\beta).$$

Proposition 2 and corollary 5 allows to bound the off-diagonal contribution, then

$$M_{f,2}(\alpha,\beta) = I_{1,1}^{D_1}(\alpha,\beta) + I_{1,1}^{D_2}(\alpha,\beta) + O\left(T^{\frac{1}{2}+\theta+\varepsilon}\right). \tag{34}$$

5.1 Diagonal contribution

We begin with a useful lemma.

Lemma 13 *The Laurent series of the meromorphic function* $s \mapsto \frac{L(f\times f,s)}{\zeta^{(N)}(2s)}$ *about* $s=1$ *can be written as*

$$\frac{L(f\times f,1+s)}{\zeta^{(N)}(2(1+s))} = \frac{\mathfrak{a}_f/2}{s} + \mathfrak{b}_f/2 + O(s).$$

Proof: We have

$$\frac{L(f\times f,1+s)}{\zeta^{(N)}(2(1+s))} = \frac{1}{\prod_{p|N}\left(1+\frac{1}{p^{1+s}}\right)}\frac{\zeta(1+s)}{\zeta(2(1+s))}L(Sym^2 f,1+s)$$

$$= \frac{N}{\nu(N)}\left(1+s\sum_{p|N}\frac{\ln p}{p+1}+O(s^2)\right)\left(L(Sym^2 f,1)+sL'(Sym^2 f,1)+O(s^2)\right)\frac{\left(1-2\frac{\zeta'(2)}{\zeta(2)}+O(s^2)\right)}{\zeta(2)}$$

$$\times\left(\frac{1}{s}+\gamma+O(s)\right).$$

An easy calculation gives the result.

□

Lemma 14 *Let* $\alpha,\beta \ll \ln T$ *be complex numbers. We have*

$$I_{1,1}^{D_1}(\alpha,\beta) = \int_{\mathbb{R}} w(t)\underset{s=0}{Res}\left[\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\frac{L(f\times f,1+\alpha+\beta+2s)}{\zeta^{(N)}(2(1+\alpha+\beta+2s))}\right]dt + O\left(T^{1/2}\right).$$

Proof: By the definition of $I_{1,1}^{D_1}(\alpha,\beta)$, we can write

$$\begin{aligned} I_{1,1}^{D_1}(\alpha,\beta) &= \sum_{m=n}\frac{\lambda_f(m)\lambda_f(n)}{m^{\frac{1}{2}+\alpha}n^{\frac{1}{2}+\beta}}\int_{\mathbb{R}} w(t)V_{\alpha,\beta}(mn,t)dt \\ &= \int_{\mathbb{R}} w(t)\frac{1}{2i\pi}\int_{(\sigma)}\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\sum_{m\geq 1}\frac{\lambda_f(m)^2}{m^{1+\alpha+\beta+2s}}dsdt. \end{aligned}$$

Since $L(f\times f,s)=\zeta^{(n)}(2s)\sum_{m\geq 1}\frac{\lambda_f(m)^2}{m^s}$, then for all positive real number σ we get

$$I_{1,1}^{D_1}(\alpha,\beta) = \int_{\mathbb{R}} w(t)\frac{1}{2i\pi}\int_{(\sigma)}\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\frac{L(f\times f,1+\alpha+\beta+2s)}{\zeta^{(N)}(2(1+\alpha+\beta+2s))}dsdt.$$

If $\alpha+\beta\neq 0$, we specialise

$$G(s) = e^{s^2}\frac{(\alpha+\beta)^2-(2s)^2}{(\alpha+\beta)^2}.$$

in order to insure $G\left(-\frac{\alpha+\beta}{2}\right)=0$. We move the integration line from $\Re s=\sigma$ to $\Re s=-A$, with $A=\frac{1}{4}+\frac{\alpha+\beta}{2}$, crossing a pole at $s=0$. Thus,

$$I_{1,1}^{D_1}(\alpha,\beta) = \int_{\mathbb{R}} w(t)\underset{s=0}{Res}\left[\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\frac{L(f\times f,1+\alpha+\beta+2s)}{\zeta^{(N)}(2(1+\alpha+\beta+2s))}\right]dt + O\left(T^{1-2A}\right).$$

□

In order to calculate the residue at $s=0$, which appears in the previous lemma, we split our proof according to the multiplicity of this pole.

Double pole case In this paragraph, we assume $\alpha+\beta=0$. Then the pole at $s=0$ in the previous lemma has multiplicity 2.

Lemma 15 *We have*

$$\operatorname*{Res}_{s=0}\left[\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\frac{L(f\times f,1+\alpha+\beta+2s)}{\zeta^{(N)}(2(1+\alpha+\beta+2s))}\right]=\frac{\mathfrak{a}_f}{2}\ln\left(\frac{t\sqrt{N}}{2\pi}\right)+\frac{\mathfrak{b}_f}{2}$$

Proof: We compute the following asymptotic behaviour at $s=0$.

$$\begin{aligned}G(s)\left(\frac{t\sqrt{N}}{2\pi}\right)^{2s}\frac{L(f\times f,1+2s)}{\zeta^{(N)}(2(1+2s))} &= \left[1+O(s^2)\right]\left[1+2s\ln\left(\frac{t\sqrt{N}}{2\pi}\right)+O(s^2)\right]\left[\frac{\mathfrak{a}_f/2}{2s}+\mathfrak{b}_f/2+O(s)\right]\\ &= \frac{\mathfrak{a}_f/2}{s}+\frac{\mathfrak{a}_f}{2}\ln\left(\frac{t\sqrt{N}}{2\pi}\right)+\frac{\mathfrak{b}_f}{2}+O(s).\end{aligned}$$

□

These results prove theorem 2 when $\alpha+\beta=0$. Precisely, by relation (34) and since $I_{1,1}^{D_1}(\alpha,\beta)=I_{1,1}^{D_2}(\alpha,\beta)$ in the considered case, the following corollary comes from lemmas 14 and 15.

Corollary 6 *Let $\alpha\ll L^{-1}$ be a complex number. For all $\varepsilon>0$, we have*

$$M_{f,2}(\alpha,-\alpha)=a_f\int_{\mathbb{R}}w(t)\ln t dt+b_f\int_{\mathbb{R}}w(t)dt+O\left(T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

Simple pole case In this paragraph, we assume $\alpha+\beta\neq 0$ and $G(s)=e^{s^2\frac{(\alpha+\beta)^2-(2s)^2}{(\alpha+\beta)^2}}$. We also set $w_2(t)=w(t)\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}$.

Lemma 16 *Let $\alpha,\beta\ll L^{-1}$ be complex numbers. For all $\varepsilon>0$, we have*

$$M_{f,2}(\alpha,\beta)=\frac{L(f\times f,1+\alpha+\beta)}{\zeta^{(N)}(2(1+\alpha+\beta))}\widehat{w}(0)+\frac{L(f\times f,1-\alpha-\beta)}{\zeta^{(N)}(2(1-\alpha-\beta))}\widehat{w}_2(0)+O\left(T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

Proof: Since the pole at $s=0$, which appears in lemma 14, is simple, we have

$$\operatorname*{Res}_{s=0}\left[\frac{G(s)}{s}g_{\alpha,\beta}(s,t)\frac{L(f\times f,1+\alpha+\beta+2s)}{\zeta^{(N)}(2(1+\alpha+\beta+2s))}\right]=\frac{L(f\times f,1+\alpha+\beta)}{\zeta^{(N)}(2(1+\alpha+\beta))}.$$

Then, by lemma 14

$$I_{1,1}^{D_1}(\alpha,\beta)=\frac{L(f\times f,1+\alpha+\beta)}{\zeta^{(N)}(2(1+\alpha+\beta))}\widehat{w}(0)+O\left(T^{1/2}\right).$$

Up to changing w by w_2, we have $I_{1,1}^{D_2}(\alpha,\beta)=I_{1,1}^{D_1}(-\beta,-\alpha)$, thus

$$I_{1,1}^{D_2}(\alpha,\beta)=\frac{L(f\times f,1-\alpha-\beta)}{\zeta^{(N)}(2(1-\alpha-\beta))}\widehat{w}_2(0)+O\left(T^{1/2}\right).$$

□

As a result, we obtain theorem 2 when $\alpha+\beta\neq 0$.

Corollary 7 *Let $\alpha,\beta\ll L^{-1}$ be complex numbers. For all $\varepsilon>0$, we have*

$$M_{f,2}(\alpha,\beta)=\mathfrak{a}_f\int_{-\infty}^{+\infty}w(t)\ln t dt+\left[\mathfrak{b}_f+\mathfrak{a}_f\ln\left(\frac{\sqrt{N}}{2\pi}\right)\right]\widehat{w}(0)+O\left(|\alpha+\beta|T(\ln T)^2+T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

Proof: Thanks to lemma 13, the previous lemma gives

$$M_{f,2}(\alpha,\beta)=\frac{\mathfrak{a}_f/2}{\alpha+\beta}\left(\widehat{w}(0)-\widehat{w}_2(0)\right)+\frac{\mathfrak{b}_f}{2}\left(\widehat{w}(0)+\widehat{w}_2(0)\right)+O\left(T|\alpha+\beta|+T^{\frac{1}{2}+\theta+\varepsilon}\right).$$

Furthermore,

$$\begin{aligned}\widehat{w}_2(0) &= \int_{\mathbb{R}}w(t)\left(\frac{t\sqrt{N}}{2\pi}\right)^{-2(\alpha+\beta)}dt=\int_{\mathbb{R}}w(t)\left(1-2(\alpha+\beta)\ln\left(\frac{t\sqrt{N}}{2\pi}\right)+O\left(|\alpha+\beta|^2\ln^2 t\right)\right)\\ &= \widehat{w}(0)-2(\alpha+\beta)\int_{\mathbb{R}}w(t)\ln\left(\frac{t\sqrt{N}}{2\pi}\right)dt+O\left(|\alpha+\beta|^2T\ln^2 T\right).\end{aligned}$$

An easy calculation gives the result.

□

We remark that, up to changing $\Delta=T/\ln T$ by $T/(\ln T)^2$, we obtain corollary 3.

5.2 Conjecture of Conrey, Farmer, Keating, Rubinstein and Snaith

We can find in [CFK$^+$05] numerous conjectures related to integral moments of L-functions. In particular, conjecture 2.5.4 predicts the asymptotic behaviour of any even integral moment of a primitive L-function on the critical line.

Conjecture 1 *Let $\mathcal{L}(s)$ be a primitive L-function. Let k be a positive integer. Then for any "suitable" weight function g, we have*

$$\int_{-\infty}^{+\infty}\left|\mathcal{L}\left(\frac{1}{2}+it\right)\right|^{2k}g(t)dt=\int_{-\infty}^{+\infty}P_k\left(w\ln\left(\frac{Q^{2/w}t}{2}\right)\right)(1+O(t^{-1/2+\varepsilon}))g(t)dt.$$

where w and Q are respectively the degree and the conductor of $\mathcal{L}$ and P_k is an explicit polynomial of degree k^2.

In this paper, we consider L-functions of holomorphic primitive cusp forms of even weight, square-free level N and trivial character. The degree of such an L-function is $w=2$ and the conductor is $Q=\sqrt{N}/\pi$ (confer equation (6)). The following conjecture is a simple rewriting of conjecture 1 in this case for the second integral moment and when $g(t)=r(t/T)$ with r a smooth function compactly supported in $[1,2]$.

Conjecture 2 *Let f be a holomorphic primitive cusp forms of even weight, square-free level N and trivial character. Then, for any $\varepsilon > 0$, we have*

$$\int_{-\infty}^{+\infty} \left| L\left(f, \frac{1}{2} + it\right)\right|^2 g(t)dt = \int_{-\infty}^{+\infty} P_1\left(2\ln\left(\frac{t\sqrt{N}}{2\pi}\right)\right) g(t)dt + O\left(T^{\frac{1}{2}+\varepsilon}\right)$$

with

$$P_1(x) = \frac{-1}{(2i\pi)^2} \oint_{|z_1|=r_1} \oint_{|z_2|=r_2} \frac{L(f\times f, 1+z_1-z_2)}{\zeta^{(N)}(2(1+z_1-z_2))} \frac{(z_2-z_1)^2}{z_1^2 z_2^2} e^{\frac{x}{2}(z_1-z_2)} dz_1 dz_2,$$

for any small positive real numbers r_1 and r_2 (ie $r_1 + r_2 < 1$).

In order to compare our corollary 2 with this conjecture, we have to compute P_1. Choosing $r_1 \neq r_2$, and since $\frac{(z_2-z_1)^2}{z_1^2 z_2^2} = \frac{1}{z_1^2} - \frac{2}{z_1 z_2} + \frac{1}{z_2^2}$, we have

$$\begin{aligned} P_1(x) &= \frac{2}{(2i\pi)^2} \oint_{|z_1|=r_1} \oint_{|z_2|=r_2} \frac{L(f\times f, 1+z_1-z_2)}{\zeta^{(N)}(2(1+z_1-z_2))} \frac{1}{z_1 z_2} e^{\frac{x}{2}(z_1-z_2)} dz_1 dz_2 \\ &= \frac{2}{2i\pi} \oint_{|z_2|=r_2} \frac{L(f\times f, 1-z_2)}{\zeta^{(N)}(2(1-z_2))} e^{-\frac{x}{2}z_2} \frac{dz_2}{z_2} \end{aligned}$$

Moreover, since

$$\frac{L(f\times f, 1-z_2)}{\zeta^{(N)}(2(1-z_2))} e^{-\frac{x}{2}z_2} = \left(\frac{-\mathfrak{a}_f/2}{z_2} + \mathfrak{b}_f + O(z_2^2)\right)\left(1 - \frac{x}{2}z_2 + O(z_2^2)\right) = \frac{-\mathfrak{a}_f/2}{z_2} + \left(\frac{\mathfrak{b}_f}{2} + \frac{x\mathfrak{a}_f}{4}\right) + O(z_2^2),$$

we obtain

$$P_1(x) = \frac{\mathfrak{a}_f}{2} x + \mathfrak{b}_f.$$

To conclude, we can see that the main terms are similar in corollary 2 and conjecture 2 and, assuming the Ramanujuan-Petersson conjecture, the error terms are also equal.

References

[BCHB85] R. Balasubramanian, J. B. Conrey, and D. R. Heath-Brown. Asymptotic mean square of the product of the Riemann zeta-function and a Dirichlet polynomial. *J. Reine Angew. Math.*, 357:161–181, 1985.

[BCY11] H. M. Bui, B. Conrey, and M. P. Young. More than 41% of the zeros of the zeta function are on the critical line. *Acta Arith.*, 150(1):35–64, 2011.

[Blo04] V. Blomer. Shifted convolution sums and subconvexity bounds for automorphic L-functions. *Int. Math. Res. Not.*, (73):3905–3926, 2004.

[Blo05] V. Blomer. Rankin-Selberg L-functions on the critical line. *Manuscripta Math.*, 117(2):111–133, 2005.

[CFK+05] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein, and N. C. Snaith. Integral moments of L-functions. *Proc. London Math. Soc. (3)*, 91(1):33–104, 2005.

[Con89] B. Conrey. More than two fifths of the zeros of the Riemann zeta function are on the critical line. *J. Reine Angew. Math.*, 399:1–26, 1989.

[DI83] J.-M. Deshouillers and H. Iwaniec. Kloosterman sums and Fourier coefficients of cusp forms. *Invent. Math.*, 70(2):219–288, 1982/83.

[Far94] D. W. Farmer. Mean value of Dirichlet series associated with holomorphic cusp forms. *J. Number Theory*, 49(2):209–245, 1994.

[Fen12] S. Feng. Zeros of the Riemann zeta function on the critical line. *J. Number Theory*, 132(4):511–542, 2012.

[Goo82] A. Good. The square mean of Dirichlet series associated with cusp forms. *Mathematika*, 29(2):278–295 (1983), 1982.

[Haf83] J. L. Hafner. Zeros on the critical line for Dirichlet series attached to certain cusp forms. *Math. Ann.*, 264(1):21–37, 1983.

[Haf87] J. L. Hafner. Zeros on the critical line for Maass wave form L-functions. *J. Reine Angew. Math.*, 377:127–158, 1987.

[Har03] G. Harcos. An additive problem in the Fourier coefficients of cusp forms. *Math. Ann.*, 326(2):347–365, 2003.

[HB79] D. R. Heath-Brown. Simple zeros of the Riemann zeta function on the critical line. *Bull. London Math. Soc.*, 11(1):17–18, 1979.

[HY10] C. P. Hughes and Matthew P. Young. The twisted fourth moment of the Riemann zeta function. *J. Reine Angew. Math.*, 641:203–236, 2010.

[IK04] H. Iwaniec and E. Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2004.

[Kim03] H. H. Kim. Functoriality for the exterior square of GL_4 and the symmetric fourth of GL_2. *J. Amer. Math. Soc.*, 16(1):139–183 (electronic), 2003. With appendix 1 by Dinakar Ramakrishnan and appendix 2 by Kim and Peter Sarnak.

[KMV02] E. Kowalski, P. Michel, and J. VanderKam. Rankin-Selberg L-functions in the level aspect. *Duke Math. J.*, 114(1):123–191, 2002.

[Lev74] N. Levinson. More than one third of zeros of Riemann's zeta-function are on $\sigma = 1/2$. *Advances in Math.*, 13:383–436, 1974.

[Ran39] R. A. Rankin. Contributions to the theory of Ramanujan's function $\tau(n)$ and similar arithmetical functions. II. The order of the Fourier coefficients of integral modular forms. *Proc. Cambridge Philos. Soc.*, 35:351–372, 1939.

[Rez10] I. S. Rezvyakova. On the zeros of Hecke L-functions and of their linear combinations on the critical line. *Dokl. Akad. Nauk*, 431(6):741–746, 2010.

[Ric06] G. Ricotta. Real zeros and size of Rankin-Selberg L-functions in the level aspect. *Duke Math. J.*, 131(2):291–350, 2006.

[Sel42] A. Selberg. On the zeros of Riemann's zeta-function. *Skr. Norske Vid. Akad. Oslo I.*, 1942(10):59, 1942.

[Ste07] J. Steuding. *Value-distribution of L-functions*, volume 1877 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.

[Ten95] G. Tenenbaum. *Introduction à la théorie analytique et probabiliste des nombres*, volume 1 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, second edition, 1995.

[Tit86] E. C. Titchmarsh. *The theory of the Riemann zeta-function*. The Clarendon Press Oxford University Press, New York, second edition, 1986. Edited and with a preface by D. R. Heath-Brown.

[You10] M. P. Young. A short proof of Levinson's theorem. *Arch. Math. (Basel)*, 95(6):539–548, 2010.

[Zha05] Q. Zhang. Integral mean values of modular L-functions. *J. Number Theory*, 115(1):100–122, 2005.

Damien BERNARD
Université Blaise Pascal
Laboratoire de Mathématiques
Campus des Cézeaux
BP 80026
63171 Aubière cedex, France
E-mail address: damien.bernard@math.univ-bpclermont.fr